



Manage resistant weeds with extended grass and broadleaf weed control for soybeans and wheat crops.

Registered crops

- Soybeans
- Spring wheat
- Winter wheat
- Field corn

Weeds controlled

- Annual sow thistle
- Barnyard grass
- Canada fleabane
- Chickweed
- Cleavers
- Common ragweed (including glyphosate-resistant biotypes)
- Crabgrass
- Dandelion
- Green foxtail (Group 1- and 2-resistant biotypes)
- Green and redroot pigweed (including Group 2- and 5-resistant biotypes)
- Lamb's-quarters (including Group 2- and triazine-resistant biotypes)
- Nightshade
- Palmer amaranth
- Ragweed
- Waterhemp (including Group 2-, 5- and glyphosate-resistant biotypes)
- Wild buckwheat
- Wild mustard
- Velvetleaf
- Volunteer canola (suppression)

Benefits

- **EXCELLENT PERFORMING RESIDUAL HERBICIDE** on tough broadleaf weeds
- **GRASSY WEED MANAGEMENT** for cleaner fields with a different herbicide mode of action
- **LONGER CONTROL** with residual activity that lasts up to 8 weeks
- **TWO ACTIVE INGREDIENTS** (proven Group 14 and 15) for management of Group 1-, 2-, 4-, 5- and glyphosate-resistant weeds
- Tank-mixable with most burndown and pre-emergent herbicides (i.e. glyphosate and BlackHawk®)
- No surfactant required

Performance tips

1. **Emerged weeds** – use with a burndown herbicide partner like glyphosate and BlackHawk
2. **Flushing weeds** – residual activity for newly emerging weeds
3. **Emerged crop** – Fierce alone can be applied up to 3 days after planting soybeans
4. Refer to label for rates and use patterns
5. DO NOT incorporate after application or prior to planting
6. Leave a small checkstrip in the field to show performance

PRE-EMERGENT RESIDUAL HERBICIDE

ACTIVE INGREDIENTS

- > Group 14 (flumioxazin)
- > Group 15 (pyroxasulfone)

TREATMENT AREA

- > One case treats 128 acres at 85 g/ac. (32 ac./jug)

APPLICATION TIMING

- > Pre-plant or pre-emergent for soybeans
- > Should be applied a minimum of 7 days prior to planting spring wheat, winter wheat and field corn (no-till only)
- > Apply up to 3 days after planting soybeans

TANK-MIX ORDER

- > Fierce
- > BlackHawk® (if used)
- > Glyphosate

NOTES

- > Planting after herbicide application requires the use of minimum disturbance planting systems
- > Fierce requires moisture to activate in the soil and provides up to 8 weeks residual control

PCP

- > 31117

1.800.868.5444 | Nufarm.ca | NufarmCA

Always read and follow label directions.

Fierce® is a registered trademark of Valent U.S.A. LLC.
BlackHawk® is a registered trademark of Nufarm Agriculture Inc.
Nufarm Agriculture Inc. is a distributor of Valent products in Canada.

73005-08/20

