



# Parasol<sup>®</sup> FL

Parasol<sup>®</sup> FL, one of the smallest and most consistent copper particle size formulations available, disperses effortlessly and completely in water providing maximum crop protection from disease.

## BENEFITS

- Easy to handle and mix
- Has a higher Metallic Copper Equivalent (MCE) than some other coppers on the market providing optimal control of tough diseases
- Superior formulation stays in suspension and quickly disperses in water
- Highly compatible with other products
- Superior rainfastness
- One of the smallest and most consistent sizes of copper particles available in Canada

## REGISTERED CROPS

- Bean
- Cucumber
- Pepper
- Potato
- Tomato

## GENERAL USAGE INFORMATION

- Use as a preventative/protective fungicide spray
- Use on a 7-14 day interval depending on disease conditions
- Adaptable for spraying with all types of equipment
- No surfactants needed
- No buffer zone required when applied as spot treatment

## TECHNICAL INFORMATION

### CHEMICAL CLASS

> M1 fungicide

### ACTIVE INGREDIENTS

> copper hydroxide 24.4%

### PACKAGING

> 2 x 10 L case

### RAINFAST

> 2 hours

### PCP #

> 25901

## SPECIFIC CROP USAGE INFORMATION

DISEASE CONTROLLED	RATE	PHI	APPLICATION INFORMATION
<b>BEAN</b>			
Bacterial blight, common ( <i>Xanthomonas campestris</i> pv. <i>phaseoli</i> )  Bacterial blight, halo ( <i>Pseudomonas syringae</i> pv. <i>phaseolicola</i> )	2.3-3.12 L/ha (0.93-1.25 L/ac.)	2	For protective sprays, apply first application when plants are 15 cm high  Apply on 7-14 day schedule depending on local conditions  Do not apply more than 6 treatments per year
<b>CUCUMBER</b>			
Angular leaf spot ( <i>Pseudomonas syringae</i> pv. <i>lachrymans</i> )	2.3-3.12 L/ha (0.93-1.25 L/ac.)	2	Apply weekly once the plants begin to vine  Do not apply more than 5 treatments per year  Minimum retreatment interval of 7 days
<b>PEPPER</b>			
Bacterial spot ( <i>Xanthomonas campestris</i> pv. <i>vesicatoria</i> and <i>X. vesicatoria</i> )	2.3-3.12 L/ha (0.93-1.25 L/ac.)	2	When disease threatens, apply on a 7-14 day interval depending on disease severity  Do not apply more than 10 treatments per year, with a minimum retreatment interval of 3 days
<b>POTATO</b>			
Early blight ( <i>Alternaria solani</i> )  Late blight ( <i>Phytophthora infestans</i> )	0.8-1.8 L/ha (0.32-0.73 L/ac.)	2	Apply at 7-10 day intervals starting when plants are 15 cm high until harvest  Combine with 1.75-2.25 kg/ha of Mancozeb  Apply 2.4 L/ha (0.97 L/ac.) at vinekill with a desiccant or alone after vinekill, prior to harvest for disease management  This late treatment may reduce infection of tubers by the late blight fungus during harvesting  Do not apply more than 10 treatments per year, with a minimum retreatment interval of 5 days
<b>TOMATO</b>			
Bacterial spot ( <i>Xanthomonas campestris</i> pv. <i>vesicatoria</i> )  Early blight ( <i>Alternaria solani</i> )  Late blight ( <i>Phytophthora infestans</i> )	2.3 L/ha (0.93 L/ac.)	2	When disease threatens, apply on a 7-10 day interval, and more frequently when disease is severe  Combine with 1.75-2.25 kg/ha Mancozeb  Do not apply more than 10 treatments per year, with a minimum retreatment interval of 3 days



1.800.868.5444 | Nufarm.ca | @NufarmCA

Always read and follow the product label for more detailed information on control of weeds, insects or disease, application directions, and use precautions. Please refer to label for more information including future label expansions that may include new crops, pests and use patterns. Please refer to label for re-entry periods.

Parasol® is a registered trademark of Nufarm Agriculture Inc. All other products are trademarks of their respective companies. 58123-01/18