

Parasol® WG

Parasol® WG provides growers with consistent and uniform copper coverage and enhanced disease protection through extremely small particle size and a high organic copper load.

BENEFITS

- Easy to handle and mix
- Eliminates clogging of sprayer and booms
- Highly compatible with other products
- Small active particle size to ensure superior coverage

REGISTERED CROPS

- Bean
- Cucumber
- Pepper
- Potato
- Sugarbeet
- Tomato

GENERAL USAGE INFORMATION

- Use as a preventative/protective fungicide spray
- Use on a 7-14 day interval depending on disease conditions
- Adaptable for spraying with all types of equipment

TECHNICAL INFORMATION

CHEMICAL CLASS

> M1 fungicide

ACTIVE INGREDIENTS

> copper hydroxide 50%

PACKAGING

> 10 kg bag

RAINFAST

> Avoid application when heavy rainfall is in the forecast

PCP

> 29063



SPECIFIC CROP USAGE INFORMATION

DISEASE CONTROLLED	RATE	PHI	APPLICATION INFORMATION
BEAN			
Bacterial blight, common (<i>Xanthomonas campestris</i> <i>pv. phaseoli</i>)	2.25-3.25 kg/ha (0.91-1.3 kg/ac.)	2	For protective sprays, apply first application when plants are 15 cm high
Bacterial blight, halo (<i>Pseudomonas syringae</i> <i>pv. phaseolicola</i>)			Apply on 7-14 day schedule depending on local conditions Do not apply more than 6 treatments per year
CUCUMBER			
Angular leaf spot (<i>Pseudomonas syringae</i> <i>pv. lachrymans</i>)	2.25-3.25 kg/ha (0.91-1.3 kg/ac.)	2	Apply weekly once the plants begin to vine Do not apply more than 5 treatments per year
PEPPER			
Bacterial spot (<i>Xanthomonas campestris</i> <i>pv.</i> <i>vesicatoria</i> and <i>X. vesicatoria</i>)	2.25-3.25 kg/ha (0.91-1.3 kg/ac.)	2	When disease threatens, apply on a 7-14 day interval depending on disease severity Do not apply more than 10 treatments per year, with a minimum retreatment interval of 3 days
POTATO			
Early blight (<i>Alternaria solani</i>)	1.1-2.5 kg/ha (.45-1.0 kg/ac.)	2	Apply at 7-10 day intervals starting when plants are 15 cm high until harvest
Late blight (<i>Phytophthora infestans</i>)			Combine with 1.75-2.25 kg/ha of Mancozeb Apply at 3.4 kg/ha at vinekill with a desiccant or alone after vinekill, prior to harvest for disease management This late treatment may reduce infection of tubers by the late blight fungus during harvesting Do not apply more than 10 treatments per year, with a minimum retreatment interval of 5 days
SUGARBEET			
Cercospora leaf spot (<i>Cercospora beticola</i>)	2.25-4.25 kg/ha (0.91-1.72 kg/ac.)	2	Start spray when disease threatens and continue for 4 to 5 applications Spray every 10-14 days depending on weather conditions at 2.25-4.25 kg/ha depending on disease severity Addition of a suitable non-herbicidal agricultural spray oil is recommended at 5.5 L/ha For specific instructions, consult local agricultural authorities Do not apply more than 6 treatments per year
TOMATO			
Bacterial spot (<i>Xanthomonas campestris</i> <i>pv.</i> <i>vesicatoria</i>)	1.75-2.25 kg/ha (0.7-0.91 kg/ac.)	2	When disease threatens, apply 2.25 kg/ha on a 7-10 day interval, more frequently when disease is severe
Early blight (<i>Alternaria solani</i>)			Combine with 1.75-2.25 kg/ha Mancozeb
Late blight (<i>Phytophthora infestans</i>)			Do not apply more than 10 treatments per year, with a minimum retreatment interval of 3 days



1.800.868.5444 | Nufarm.ca

Always read and follow the product label for more detailed information on control of weeds, insects or disease, application directions, and use precautions. Please refer to label for more information including future label expansions that may include new crops, pests and use patterns. Please refer to label for re-entry periods.

Parasol® is a registered trademark of Nufarm Agriculture Inc. PRO-CERT ORGANIC is a trademark of PRO-CERT ORGANIC SYSTEMS, LTD.
51481-0117