

# Mextrol® 450

## MCPA + Bromoxynil

### TECHNICAL INFORMATION

<b>Chemical class:</b>	Group 4 + 6
<b>Application rate:</b>	1.25 L/ha (0.5 L/ac.)
<b>Re-entry interval:</b>	24 hours in most uses
<b>Packaging:</b>	2 x 10L case

### REGISTERED CROPS

- Underseeded winter wheat
- Wheat – Spring, winter, durum
- Oats
- Barley
- Rye
- Flax
- Corn (field, sweet)
- Canary Seed
- Seedling and established grasses

### SPECIFIC USE INSTRUCTIONS

- Apply Mextrol to young, actively growing weeds
- For underseeded crop apply when red clover is in the 1-3 trifoliate stage and wheat provides a protective canopy over the clover
- Apply to cereals in the 2-leaf to flag-leaf stage
- Follow specific label instructions for stage and applications when tank-mixed with other herbicides
- One 10L jug treats 8 ha (20 ac)

### BENEFITS

- Dual mode of action
- Outstanding Wild buckwheat control
- Gentle on underseeded red clover
- Tank-mixable with Boost, Refine SG and most grass herbicides
- Excellent partner with foliar fungicides and fertilizers

### WEEDS CONTROLLED

- Bluebur
- Canada Thistle (top growth)
- Cocklebur
- Common ragweed
- Common groundsel
- Cow cockle
- Flixweed
- Kochia
- Lady's-thumb
- Lamb's-quarters
- Mustard species
- Night flowering catchfly
- Perennial sow-thistle (top growth)
- Prickly lettuce (suppression)
- Redroot pigweed
- Russian Thistle
- Scentless chamomile
- Shepherd's-purse
- Smartweeds
- Stinkweed
- Volunteer sunflower
- Volunteer rapeseed (including canola)
- Velvetleaf
- Wild buckwheat (and buckwheat species)

1.800.868.5444

Nufarm.ca

Mextrol® is a registered trademark of Nufarm Agriculture Inc. Always read and follow label directions.



Nufarm

Grow a better tomorrow.