

BlackHawk® pre-seed herbicide enters the heat of the battle with fast, flexible and complete broadleaf weed control ahead of cereals, soybeans and corn.

Registered crops

PRE/POST-SEED APPLICATIONS

- Barley
- Canary seed
- Field corn
- Oats
- Rye (spring, fall)
- Soybeans
- Summerfallow
- Triticale
- Wheat (spring, durum, winter)

FALL APPLICATIONS

- All crops

Weeds controlled with a glyphosate tank-mix

- Canada thistle
- Cleavers (including glyphosate-resistant biotypes)
- Cocklebur
- Cow cockle (suppression)
- Dandelion (spring seedling and season long suppression)
- Flixweed
- Kochia (including Group 2-, 4- and glyphosate-resistant biotypes)
- Lamb's-quarters (including Group 2- and 5-resistant biotypes)
- Mustards (except dog and tansy)
- Narrow-leaved hawk's-beard (including Group 2-resistant biotypes)
- Night-flowering catchfly
- Pigweed (redroot and Russian)
- Prickly lettuce
- Russian thistle
- Shepherd's-purse
- Sow thistle
- Stinkweed
- Sweet clover
- Volunteer canola (including all herbicide-resistant biotypes)
- Wild buckwheat (suppression)
- Wild radish

Benefits

- **PROVEN RESISTANCE MANAGEMENT** with two non-Group 2 modes of action that target broadleaf weeds
- **COMPLETE CONTROL** through systemic and contact activity
- **BETTER PERFORMANCE** compared to glyphosate alone for fall or pre-seed burndown
- Makes in-crop weed control easier and more effective by having only new, small weeds present at time of application
- Provides broad-spectrum weed control, including excellent control of narrow-leaved hawk's-beard in the fall and at 1-2 leaf in spring
- No surfactant needed when applied with glyphosate

ACTIVE INGREDIENTS

- > Group 14 (pyraflufen-ethyl)
- > Group 4 (2,4-D ester)

RAINFAST

- > 1 hour

TREATMENT AREA

- > One case treats 60 acres at 300 mL/ac.
- > One drum treats 320 acres at 300 mL/ac.
- > One tote treats 1,600 acres at 300 mL/ac. (contact local retailer to special order)

TANK-MIX ORDER

- > BlackHawk
- > Glyphosate

NOTES

- > Fall applications can be made until freeze up with no cropping restrictions the following season
- > In the spring, applications can be made after seeding cereals prior to emergence except for oats (apply a minimum of 7 days prior to seeding)
- > Can be applied up to 3 days after seeding soybeans (ensure soybeans are seeded minimum 1 inch deep)
- > For best results, use higher water volumes for good coverage and contact activity