



A pre-emergent soil active herbicide that offers extended activity on emerging spring broadleaf and grassy weeds for up to 8 weeks after activation.

Registered crops

- Chickpeas
- Field corn
- Field peas
- Lentils (fall application prior to large green and small red varieties)
- Soybeans
- Spring wheat (not including durum)
- Winter wheat

Weeds controlled

- Annual sow-thistle
- Barnyard grass
- Chickweed
- Cleavers
- Dandelion
- Downy brome (suppression)
- Green foxtail
- Green and redroot pigweed (including Group 2- and 5-resistant biotypes)
- Japanese brome (suppression)
- Kochia (including Group 2-, 4- and glyphosate-resistant biotypes)
- Lamb's-quarters (including Group 2- and triazine-resistant biotypes)
- Russian thistle (suppression)
- Wild buckwheat
- Wild mustard
- Wild oats (suppression)
- Volunteer canola (suppression)

Benefits

- **BETTER PERFORMANCE** than other leading soil active herbicides on tough broadleaf weeds
- **ACTIVITY ON GRASSY WEEDS** with a different mode of action to manage Group 1- and 2-resistant biotypes
- **EXTENDED CONTROL** with activity that lasts up to 8 weeks
- **TWO ACTIVE INGREDIENTS** (proven Group 14 and 15) for management of Group 1-, 2-, 4-, 5- and glyphosate-resistant weeds
- Convenient application eliminates the need for granular broadcast herbicide products
- Tank-mixable with most burndown and pre-emergent herbicides (i.e. glyphosate, BlackHawk® and/or GoldWing®)

Performance tips

1. **Emerged weeds** – use with a burndown herbicide partner like glyphosate and BlackHawk or GoldWing
2. **Flushing weeds** – residual activity for newly emerging weeds
3. **Emerged crop** – Fierce alone can be applied up to 3 days post seeding soybeans, chickpeas and field peas
4. Refer to label for rates and use patterns
5. DO NOT incorporate after application or prior to seeding
6. Pre-seeding applications require the use of minimum disturbance seeding systems
7. Non-ionic surfactant required if glyphosate is not being tank-mixed
8. Leave a small checkstrip in the field to show performance

ACTIVE INGREDIENTS

- > Group 14 (flumioxazin)
- > Group 15 (pyroxasulfone)

TREATMENT AREA

- > One case treats 128 acres at 85 g/ac. (32 ac./jug)

APPLICATION TIMING

- > Fall apply as late as possible but prior to freeze up for any registered crop
- > Pre-plant or pre-emergent for soybeans, chickpeas and field peas
- > Should be applied a minimum of 7 days prior to seeding field corn, spring wheat and winter wheat
- > Apply up to 3 days after seeding soybeans, chickpeas and field peas

TANK-MIX ORDER

- > Fierce
- > BlackHawk® or GoldWing® (if used)
- > Glyphosate

NOTES

- > Seeding after application requires the use of minimum to zero till drills
- > Fierce requires moisture to activate in the soil

1.800.868.5444 | Nufarm.ca |  NufarmCA

Always read and follow label directions.

Fierce® is a registered trademark of Valent U.S.A. LLC. BlackHawk® and GoldWing are registered trademarks of Nufarm Agriculture Inc. Nufarm Agriculture Inc. is a distributor of Valent products in Canada.

73005-08/20



 **Nufarm**
Grow a better tomorrow