

Statue™ herbicide is an effective post-emergent grass herbicide registered on fruit and vegetable crops.

Statue™

Benefits

- Superior over-the-top control of both annual and perennial grasses
- Excellent plant safety allows for most crops to be treated at all stages so that applications can be targeted to the optimum weed stage
- Systemic action results in exceptional weed control
- Tank-mix flexibility with several herbicide partners
- One case will treat 40-80 ac. of horticultural crop

Registered crops

- Carrot
- Blueberry (highbush)
- Dry onion
- Potato
- Spinach
- Sunflower
- Red garden (table) beet
- Parsnip
- Radish

General usage information

- Statue herbicide must be sprayed with the adjuvant Carrier
- Thorough coverage of the foliage is important for consistent grass control
- Product will be less effective when plants are stressed by lack of moisture, excessive moisture, low temperature and/or low humidity
- Always adhere to the pre-harvest interval for each crop
- Do not use in greenhouses



Technical information

HERBICIDE GROUP

> Group 1

ACTIVE INGREDIENTS

> clethodim 240 g/L

PACKAGING

> 2 x 3 L case (requires matching case of Carrier)

RAINFAST

> 1 hour

PCP

> 32885

> Carrier PCP # 30639

RATE PHI APPLICATION INFORMATION

DRY ONION

0.38 L/ha (0.15 L/ac.)
plus 0.5% v/v of Carrier

45

Apply a maximum of 1 application per year
Apply post emergent when the crop is in the 1-4 leaf stage

POTATO

0.19 L/ha (0.08 L/ac.) to 0.38 L/ha (0.15 L/ac.)
plus 0.5% v/v of Carrier

60

Ground and aerial application

SPINACH

0.19 L/ha (0.08 L/ac.)
plus 0.5% v/v of Carrier

14

Apply a maximum of 2 applications per year. If repeat application is required, allow at least 14 days between first and second application. Apply post emergent by ground application when grassy weeds are between the 2-6 leaf stage

SUNFLOWER

0.38 L/ha (0.15 L/ac.)
plus 0.5% v/v of Carrier

72

Ground and aerial application

CARROT, PARSNIP, RADISH, RED GARDEN (table) BEET

0.19 L/ha (0.08 L/ac.) to 0.38 L/ha (0.15 L/ac.)
plus 0.5% v/v of Carrier

30

Apply a maximum of 2 applications per year. If repeat application is required, allow at least 14 days between first and second application. Do not apply more than 0.38 L/ha (0.15 L/ac.) per crop season. Apply in a minimum spray volume of 110 L/ha



1.800.868.5444 | Nufarm.ca | NufarmHortCA