



A strong pre-seed burndown solution for battling tough broadleaf weeds in cereals, soybeans and corn.

### Registered crops

#### PRIOR TO SEEDING

- Barley
- Canary seed
- Field corn
- Oats
- Rye (spring, fall)
- Soybeans
- Summerfallow
- Triticale
- Wheat (spring, durum, winter)

#### FALL APPLICATIONS

- All crops

### Weeds controlled with a glyphosate tank-mix

- Canada thistle
- Cleavers (including glyphosate-resistant biotypes)
- Cocklebur
- Cow cockle (suppression)
- Dandelion (spring seedling and season long suppression)
- Flixweed
- Kochia (including Group 2-, 4- and glyphosate-resistant biotypes)
- Lamb's-quarters (including Group 2- and 5-resistant biotypes)
- Mustards (except dog and tansy)
- Narrow-leaved hawk's-beard (including Group 2-resistant biotypes)
- Night-flowering catchfly
- Pigweed (redroot and Russian)
- Prickly lettuce
- Russian thistle
- Shepherd's-purse
- Sow thistle
- Stinkweed
- Sweet clover
- Volunteer canola (including all herbicide-resistant biotypes)
- Wild buckwheat (suppression)\*
- Wild radish

### Benefits

- **PROVEN RESISTANCE MANAGEMENT** with two non-Group 2 modes of action that target broadleaf weeds
- **FASTER COMPLETE CONTROL** through systemic and contact activity
- **BETTER PERFORMANCE** compared to glyphosate alone for fall or pre-seed burndown
- Makes in-crop weed control easier and more effective by having only new, small weeds present at time of application
- Provides broad-spectrum weed control, including excellent control of narrow-leaved hawk's-beard in the fall and at 1-2 leaf in spring
- No surfactant needed when applied with glyphosate

\*for optimum control of wild buckwheat, use a rate of 445 mL/ac.

### ACTIVE INGREDIENTS

- > Group 14 (pyraflufen-ethyl)
- > Group 4 (2,4-D ester)

### RAINFAST

- > 1 hour

### TREATMENT AREA

- > One case treats 60 acres at 300 mL/ac.
- > One drum treats 320 acres at 300 mL/ac.
- > One tote treats 1,600 acres at 300 mL/ac. (contact local retailer to special order)

### TANK-MIX ORDER

- > BlackHawk
- > Glyphosate

### NOTES

- > Fall applications can be made until freeze up with no cropping restrictions the following season
- > Cereals can be seeded immediately following application (except oats)
- > Can be applied up to 3 days after seeding soybeans (ensure soybeans are seeded minimum 1 inch deep)
- > For best results, use higher water volumes for good coverage and contact activity

**REAL FARM** REWARDS

1.800.868.5444 | Nufarm.ca | NufarmCA

