



Manage resistant weeds with extended grass and broadleaf weed control in soybeans and wheat.

Registered crops

- Field corn
- Soybeans
- Spring wheat (not including durum)

Weeds controlled

- Chickweed
- Dandelion
- Green foxtail
- Green and redroot pigweed (including Group 2- and 5-resistant biotypes)
- Kochia (including Group 2-, 4- and glyphosate-resistant biotypes)
- Lamb's-quarters (including Group 2- and triazine-resistant biotypes)
- Wild buckwheat
- Wild mustard
- Volunteer canola (suppression)

Benefits

- **BETTER PERFORMANCE** than other leading residual herbicides on tough broadleaf weeds
- **GRASSY WEED MANAGEMENT** for cleaner fields with a different mode of action
- **LONGER CONTROL** with residual activity that lasts up to 8 weeks
- **TWO ACTIVE INGREDIENTS** (proven Group 14 and a new Group 15) for management of Group 1-, 2-, 4-, 5- and glyphosate-resistant weeds
- Convenient application eliminates the need for granular broadcast herbicide products
- Tank-mixable with most burndown and pre-emergent herbicides (i.e. glyphosate, BlackHawk and/or GoldWing)

Performance tips

1. Emerged weeds – use with a burndown herbicide partner like glyphosate and BlackHawk or GoldWing
2. Flushing weeds – residual activity for newly emerging weeds
3. Emerged crop – Fierce alone can be used up to 3 days post seeding soybeans
4. Refer to label for rates and use patterns
5. DO NOT incorporate after application or prior to seeding
6. Pre-seeding applications require the use of minimum disturbance seeding systems
7. Non-ionic surfactant required if glyphosate is not being tank-mixed
8. Leave a small checkstrip in the field to show performance

PRE-EMERGENT RESIDUAL HERBICIDE

ACTIVE INGREDIENTS

- > Group 14 (flumioxazin)
- > Group 15 (pyroxasulfone)

TREATMENT AREA

- > One case treats 128 acres at 85 g/ac. (32 ac./jug)

APPLICATION TIMING

- > Fall apply as late as possible but prior to freeze up for any registered crop
- > Pre-plant or pre-emergent for soybeans
- > Should be applied a minimum of 7 days prior to seeding field corn and spring wheat
- > Apply up to 3 days after seeding soybeans

TANK-MIX ORDER

- > Fierce
- > BlackHawk or GoldWing (if used)
- > Glyphosate

NOTES

- > Seeding after application requires the use of minimum to zero till drills
- > Fierce requires moisture to activate in the soil

REAL FARM REWARDS™

1.800.868.5444 | Nufarm.ca | [NufarmCA](https://twitter.com/NufarmCA)

Always read and follow label directions.
Fierce® is a registered trademark of Valent U.S.A. LLC. Nufarm Agriculture Inc. is a distributor of Valent products in Canada.
Real Farm Rewards™ is a trademark of the Bayer Group, Monsanto Canada ULC licensee.

62818-01/19



 **Nufarm**
Grow a better tomorrow