



A strong burndown solution for battling tough broadleaf weeds in soybeans, cereals and corn.

### Registered crops

- Barley
- Field corn
- Oats
- Rye
- Soybeans
- Wheat (spring, winter, durum)

### Weeds controlled with a glyphosate tank-mix

- Burdock (prior to 4 leaf)
- Canada fleabane (suppression) (including glyphosate-resistant)
- Cocklebur
- Common ragweed
- Daisy fleabane
- Dandelion
- Giant ragweed (including glyphosate-resistant)
- Lamb's-quarters
- Mallow
- Mustards
- Night-flowering catchfly
- Pigweeds
- Plantain
- Prickly lettuce
- Shepherd's-purse
- Sow thistle
- Stinkweed
- Wild buckwheat
- Volunteer canola

### Benefits

- **THREE MODES OF ACTION** when tank-mixed with glyphosate
- **FLEXIBILITY** to apply up to one day post-planting in soybeans
- **STRONGER RESISTANCE MANAGEMENT** with two active ingredients that control glyphosate-, Group 2- and triazine-resistant biotypes
- **FASTER CONTROL** through systemic and contact activity for broad-spectrum weed control
- **WIDE SPECTRUM CONTROL** of broadleaf weeds
- Convenient all-in-one formulation
- Tank-mixable with most pre-emergent herbicides
- No surfactant required

### ACTIVE INGREDIENTS

- > Group 14 (pyraflufen-ethyl)
- > Group 4 (2,4-D ester)

### RAINFAST

- > 1 hour

### TREATMENT AREA

- > One case treats 40 acres at 445 mL/ac. (1.1 L/ha)

### TANK-MIX ORDER

- > BlackHawk
- > Glyphosate

### NOTES

- > For best results, use higher water volumes for good coverage and contact activity
- > Apply to actively growing weeds

### PCP #

- > 32111