



A pre-emergent residual herbicide for early season broadleaf weed control in soybeans, wheat and field corn.

Registered crops

- Field corn
- Soybeans
- Wheat

Weeds controlled

- Canada fleabane
- Chickweed
- Common ragweed
- Dandelion (seedling)
- Green foxtail (suppression)
- Kochia
- Lamb's-quarters
- Nightshade
- Palmer amaranth
- Pigweeds
- Volunteer canola – all biotypes (suppression)
- Waterhemp

Benefits

- **EARLY SEASON CONTROL** eliminates broadleaf weed pressure early on in the crop
- **EXTENDED, RESIDUAL CONTROL** (up to 8 weeks) on tough weeds
- **RESISTANCE MANAGEMENT TOOL** helps to control glyphosate-resistant weeds
- Apply in fall or spring (check label for specific crop registrations)
- Extended early season control for extra in-crop herbicide flexibility and to improve the efficacy of in-crop herbicide applications by having to target fewer, smaller weeds
- Yield potential is increased through early elimination of weed competition
- More re-cropping flexibility the following year
- Tank-mixable with most burndown and pre-emergent herbicides (i.e. glyphosate and BlackHawk®)

Performance tips

1. **Emerged weeds** – use with glyphosate
2. **Flushing weeds** – residual activity for newly emerging weeds
3. **Emerged crop** – Valtera alone can be used up to 3 days post planting but prior to emergence of registered crops other than wheat
4. Refer to label for rates and use patterns
5. **DO NOT** incorporate after application or prior to planting

PRE-EMERGENT RESIDUAL HERBICIDE

ACTIVE INGREDIENT

- > Group 14 (flumioxazin)

TREATMENT AREA

- > One case treats 106-160 acres at 56-85 g/ac. rate

APPLICATION

- > Tank-mix with glyphosate and/or BlackHawk for enhanced control of emerged weeds

APPLICATION TIMING

- > Pre-plant or pre-emergent for soybeans
- > Should be applied a minimum of 7 days prior to planting wheat and field corn
- > Apply Valtera up to 3 days after seeding soybeans

TANK-MIX ORDER

- > Valtera
- > BlackHawk (if used)
- > Glyphosate

NOTES

- > Planting after application requires the use of minimum to zero till drills
- > Moisture is necessary to activate Valtera in the soil for residual weed control

PCP

- > 29230

1.800.868.5444 | Nufarm.ca | NufarmCA

Always read and follow label directions.

Valtera™ is a trademark of Valent U.S.A. LLC. BlackHawk® is a registered trademark of Nufarm Agriculture Inc. Nufarm Agriculture Inc. is a distributor of Valent products in Canada.

63013-01/19

