

FIGHT GREY MOULD WITH BIOLOGICAL POWER.

Protect your greenhouse and outdoor ornamentals from emergence to harvest with Botector® Biological Fungicide. Botector suppresses grey mould on many crops including chrysanthemums, hydrangeas, poinsettias, and roses.

KEY BENEFITS

- Protects 24 ornamental crops
- Delivers effective preventive control of grey mould under low to moderate disease pressure
- May be applied weekly as needed
- Labelled for use throughout the entire growing season (from emergence/transplant up to one week prior to harvest)
- The active ingredient, *Aureobasidium pullulans*, may be compatible with certain commonly used pesticides
- May be applied as a foliar spray alone or in alternating spray programs with other registered crop protection products

DISEASES CONTROLLED	APPLICATION RATES	APPLICATION NOTES
grey mould (suppression)	1 kg/ha in 500-2000 L/ha water volume	Begin preventive applications soon after emergence or transplant if climate conditions are favourable for infection as well as at first sign of disease onset. Repeat as needed on a 7-day interval up to harvest. Preferably apply in the morning or evening hours and during cool temperatures.

DIRECTIONS FOR USE

Clean tank before using. Botector must be diluted in water. Keep the water temperature below 25°C. Fill the spray tank partially with clean water and begin agitation. Add the specific amount of Botector. To obtain a proper spray concentration, the tank has to be filled up to the necessary volume. The mixture should be stirred during application. Already mixed spray solution should not stand overnight or for more than eight hours to assure viability of *Aureobasidium pullulans*. Do not store unused product in the tank.

- Application of Botector prior to the disease onset or in the early stages of the disease development provides the best results for management of the target plant disease.
- Use the specified maximum label rates and shorter spray intervals for conditions conducive to rapid disease development. For proper application, determine the surface area to be treated, the label use rate and select appropriate water amount to ensure good canopy penetration and coverage of plant parts to be protected.
- Spray plants to the point of running-off.
- Prepare only the amount of spray solution required to treat the measured acreage.

See label for complete directions for use.

PCP NO.	31248	FRAC NO.	Not Classified
ACTIVE INGR.	<i>Aureobasidium pullulans</i> (DSM 14940 and DSM 14941)		
FORMULATION	wetable granule		
CHEM. FAMILY	biological		
PACKAGE SIZE	10 x 1.2 kg jug		
RAINFAST	avoid application when heavy rain is forecast		
SURFACTANT NEEDED	no		
KEY USES	outdoor- and greenhouse-grown ornamentals		



1.800.868.5444 • nufarm.ca

For specific application rates, directions, mixing instructions and precautions, read the product label. Please visit www.nufarm.ca to download a full label.