

Valtera™ EZ

NEW

A pre-emergent residual herbicide for early season broadleaf weed control in soybeans, wheat and field corn. Now available in a convenient liquid formulation.

Registered crops

- Field corn
- Soybeans
- Wheat

Weeds controlled

- Canada fleabane
- Chickweed
- Common ragweed
- Dandelion (seedling)
- Green foxtail (suppression)
- Lamb's-quarters
- Nightshade
- Palmer amaranth
- Pigweeds
- Volunteer canola – all biotypes (suppression)
- Waterhemp (including Group 2-, 5- and glyphosate-resistant biotypes)

Benefits

- **EARLY SEASON CONTROL** eliminates broadleaf weed pressure early in the growing season
- **EXTENDED, RESIDUAL CONTROL** (up to 8 weeks) on tough weeds
- **RESISTANCE MANAGEMENT TOOL** helps to control glyphosate-resistant weeds
- Apply in fall or spring (check label for specific crop registrations)
- Extended early season weed control for extra in-crop herbicide flexibility and to improve the efficacy of in-crop herbicide applications by having to target fewer, smaller weeds
- Yield potential is increased through early elimination of weed competition
- Re-cropping flexibility the following year
- Tank-mixable with most burndown and pre-emergent herbicides (i.e. glyphosate and BlackHawk®)

Performance tips

1. **Emerged weeds** – use with a burndown herbicide partner like glyphosate and BlackHawk
2. **Flushing weeds** – residual activity for newly emerging weeds
3. **Emerged crop** – Valtera EZ alone can be used up to 3 days post planting, but prior to emergence of soybeans only
4. Refer to label for rates and use patterns
5. DO NOT incorporate after application or prior to planting
6. Pre-plant applications require the use of minimum disturbance planting systems
7. Leave a small checkstrip in the field to show performance

PRE-EMERGENT RESIDUAL HERBICIDE

ACTIVE INGREDIENT

- > Group 14 (flumioxazin)

TREATMENT AREA

- > One case treats 106-160 acres at 60-90 mL/ac. rate

APPLICATION

- > Tank-mix with glyphosate and BlackHawk for enhanced control of emerged weeds

APPLICATION TIMING

- > Pre-plant or pre-emergent for soybeans
- > Should be applied a minimum of 7 days prior to planting wheat and field corn
- > Apply Valtera EZ up to 3 days after planting soybeans

TANK-MIX ORDER

- > Valtera EZ
- > BlackHawk (if used)
- > Glyphosate

NOTES

- > Planting after application requires the use of minimum disturbance planting systems
- > Moisture is necessary to activate Valtera EZ in the soil for residual weed control

PCP

- > 33523

1.800.868.5444 | Nufarm.ca |  NufarmCA

Always read and follow label directions.

Valtera™ is a trademark of Valent U.S.A. LLC. BlackHawk® is a registered trademark of Nufarm Agriculture Inc. Nufarm Agriculture Inc. is a distributor of Valent products in Canada.

67389-01/20



 **Nufarm**
Grow a better tomorrow