

Valtera™ EZ

A pre-emergent soil active herbicide for early season broadleaf weed control in pulses, wheat and soybeans. Now available in a convenient liquid formulation.

Registered crops

- Chickpeas
- Field corn
- Field peas
- Lentils (large green and small red varieties)
- Soybeans
- Wheat

Weeds controlled

- Canada fleabane*
- Chickweed
- Dandelion (seedling)
- Green foxtail (suppression)
- Kochia (including Group 2-, 4- and glyphosate-resistant biotypes)
- Lamb's-quarters
- Palmer amaranth*
- Pigweeds
- Volunteer canola – all biotypes (suppression)
- Waterhemp*

*Farmers in southern Manitoba should watch for these weeds moving northward from the US.

Benefits

- **EARLY SEASON CONTROL** helps eliminate broadleaf weed pressure in pulses, wheat and soybeans during critical weed periods
- **EXTENDED CONTROL** with soil activity that provides longer lasting control of tough weeds (up to 8 weeks)
- **RESISTANCE MANAGEMENT TOOL** that helps manage weeds on the Valtera label that are resistant to glyphosate, Group 2-, and Group 4- herbicides
- **RE-CROPPING FLEXIBILITY** – most crops not listed on the label can be seeded 12 months after application
- Apply in fall or spring (check label for specific crop registrations)
- No hassle application that doesn't include the handling and incorporation of bulky granular herbicide products
- Extended early season control for extra in-crop herbicide flexibility that helps to improve the efficacy of in-crop herbicide applications by targeting fewer, smaller weeds
- Yield potential is increased through early elimination of weed competition
- Tank-mixable with most burndown and pre-emergent herbicides (i.e. glyphosate, GoldWing and/or BlackHawk)

Performance tips

1. **Emerging weeds** – use with a burndown herbicide partner like glyphosate and BlackHawk or GoldWing
2. **Flushing weeds** – residual activity for newly emerging weeds
3. **Emerging crop** – Valtera EZ alone can be used up to 3 days post seeding, but prior to emergence of soybeans, chickpeas and field peas
4. Refer to label for rates and use patterns
5. DO NOT incorporate after application or prior to seeding
6. Requires the use of minimum disturbance seeding systems
7. Leave a small checkstrip in the field to show performance

PRE-EMERGENT SOIL ACTIVE HERBICIDE

ACTIVE INGREDIENT

- > Group 14 (flumioxazin)

TREATMENT AREA

- > One case treats 160 acres at 60 mL/ac.

APPLICATION

- > Tank-mix with glyphosate and/or BlackHawk®/GoldWing® for enhanced control of emerged weeds

APPLICATION TIMING

- > Fall apply as late as possible but prior to freeze up for any registered crop
- > Pre-plant or pre-emergent for soybeans, field peas and chickpeas
- > Should be applied a minimum of 7 days prior to seeding field corn, wheat and lentils
- > Apply up to 3 days after seeding soybeans, chickpeas and field peas
- > Must be applied 30 days prior to seeding durum wheat

TANK-MIX ORDER


- > Valtera EZ
- > BlackHawk or GoldWing (if used)
- > Glyphosate

NOTES

- > Seeding after application requires the use of minimum to zero till drills
- > Valtera requires moisture to activate in the soil



MIX IT UP™

Brought to you by 

Save
\$2/acre

1.800.868.5444 | Nufarm.ca |  NufarmCA

Always read and follow label directions.

Valtera™ is a trademark of Valent U.S.A. LLC. Nufarm Agriculture Inc. is a distributor of Valent products in Canada. Bayer®, Bayer Cross, BayerValue™ and Mix it Up™ are the trademarks of the Bayer Group or its licensors. BlackHawk® and GoldWing® are registered trademarks of Nufarm Agriculture Inc.



66601-04/20

 **Nufarm**
Grow a better tomorrow