

Valtera™ EZ

A pre-emergent residual herbicide for early season broadleaf weed control in soybeans, wheat and field corn. Now available in a convenient liquid formulation.

Registered crops

- Field corn
- Soybeans
- Wheat (spring, winter)

Weeds controlled

- Annual sow-thistle
- Canada fleabane
- Chickweed
- Cleavers (suppression)
- Common ragweed
- Dandelion (seedling)
- Green foxtail (suppression)
- Lamb's-quarters
- Nightshade
- Palmer amaranth
- Pigweeds
- Volunteer canola (suppression including all herbicide-tolerant biotypes)
- Waterhemp (including Group 2-, 5- and glyphosate-resistant biotypes)
- Wild buckwheat (suppression)

Benefits

- **EARLY SEASON CONTROL** eliminates broadleaf weed pressure early in the growing season
- **EXTENDED, RESIDUAL CONTROL** (up to 8 weeks) on tough weeds
- **RESISTANCE MANAGEMENT TOOL** helps to control glyphosate-resistant weeds
- Apply in fall or spring (check product label for specific crop registrations)
- Extended early season weed control for extra in-crop herbicide flexibility and to improve the efficacy of in-crop herbicide applications by having to target fewer, smaller weeds
- Yield potential is increased through early elimination of weed competition
- Re-cropping flexibility the following year
- Tank-mixable with most burndown and pre-emergent herbicides (i.e. glyphosate and BlackHawk®)

Performance tips

1. **Emerged weeds** – use with a burndown herbicide partner like glyphosate and BlackHawk
2. **Flushing weeds** – residual activity for newly emerging weeds
3. **Application flexibility** – Valtera EZ alone can be used up to 3 days post planting, but prior to emergence of soybeans only
4. Refer to product label for rates and use patterns
5. **DO NOT** incorporate after application or prior to planting
6. Leave a small checkstrip in the field to show performance

ACTIVE INGREDIENT

- > Group 14 (flumioxazin)

TREATMENT AREA

- > One case treats 106-160 acres at 60-90 mL/ac.

APPLICATION

- > Tank-mix with glyphosate and BlackHawk for enhanced control of emerged weeds

APPLICATION TIMING

- > Pre-emergent for soybeans
- > Should be applied a minimum of 7 days prior to planting wheat and field corn
- > Apply Valtera EZ up to 3 days after planting soybeans

TANK-MIX ORDER

- > Valtera EZ
- > BlackHawk (if used)
- > Glyphosate

NOTES

- > Planting after application requires the use of minimum disturbance planting systems
- > Moisture is necessary to activate Valtera EZ in the soil for residual weed control

PCP

- > 33523

1.800.868.5444 | Nufarm.ca | [NufarmCA](https://twitter.com/NufarmCA)

Always read and follow label directions.

Valtera™ is a trademark of Valent U.S.A. LLC.
BlackHawk® is a registered trademark of Nufarm Agriculture Inc.
Nufarm Agriculture Inc. is a distributor of Valent Canada Inc. products in Canada.
75746-0221



 **Nufarm**
Grow a better tomorrow