

Quash® SC

Helps manage the impact of sclerotinia in canola, and powdery mildew in pulses, as well as the suppression of many other diseases.

REGISTERED CROPS

- Canola
- Pulses (field peas, chickpeas, lentils)
- Dry beans
- Potatoes
- Sunflowers

DISEASES CONTROLLED

Canola

- Sclerotinia stem rot (*Sclerotinia sclerotiorum*)

Field Peas

- Powdery mildew (*Erysiphe pisi*)

Potatoes

- Early blight (*Alternaria solani*)

Sunflowers

- Rust (*Puccinia helianthi*)

See the product label for additional diseases suppressed in chickpeas, lentils, dry beans and potatoes.

BENEFITS

- **EXCELLENT CONTROL OF SCLEROTINIA** in canola and powdery mildew in field peas
- **SYSTEMIC FUNGICIDE** that is absorbed quickly through the cuticle of the plant
- **PROVIDES RESIDUAL ACTIVITY**, and moves outward in the leaf tissue from the initial point of contact within the plant
- No adjuvant required to achieve maximum performance

For specific crop usage information see label for details.

ACTIVE INGREDIENT

- Group 3 (metconazole)

RAINFAST

- 2 hours

TREATMENT AREA

- One case (2 x 4.8 L) treats 160 acres

CROP STAGING

- Canola: Apply between 20%–50% bloom
- Pulses: Make first application at 20–50% bloom before disease symptoms are visible

NOTES

- Apply prior to disease development
- For best results apply with 20 gal./ac. of water by ground or 5 gal./ac. by air