

# Ransack®

Ransack® is a broad-spectrum in-crop herbicide for field peas, soybeans, faba beans and dry beans. The convenient all-in-one formulation also delivers re-cropping flexibility.

## REGISTERED CROPS

- Field peas
- Dry beans
- Faba beans
- Soybeans
- Alfalfa, clover and sainfoin for seed production (see label)

## WEEDS CONTROLLED (WITH UAN ADDED)

- Barnyard grass
- Cleavers (including Group 2-resistant biotypes)
- Cow cockle
- Green and yellow foxtail
- Green smartweed
- Hemp nettle (suppression)
- Japanese brome (suppression)
- Kochia (suppression)
- Lamb's-quarters
- Persian darnel
- Redroot pigweed
- Round-leaved mallow (suppression)
- Russian thistle
- Shepherd's purse
- Sowthistle (annual and perennial) (suppression)
- Stinkweed
- Volunteer canola (including all herbicide-tolerant biotypes)
- Volunteer cereals (barley, canary seed, durum wheat, spring wheat, oats)
- Volunteer lentils (including all herbicide-tolerant biotypes)
- Wild buckwheat (suppression)
- Wild mustard (including Group 2-resistant)
- Wild oats

## BENEFITS

- Re-cropping flexibility
- Multiple modes of action to manage herbicide-resistant weeds
- Convenient all-in-one formulation
- No surfactant required

**For specific crop usage information see label for details.**

## ACTIVE INGREDIENTS

- Group 2 (imazamox)
- Group 6 (bentazon)

## RAINFAST

- 6 hours

## TREATMENT AREA

- One case treats 40 acres
- One drum treats 240 acres (contact local retailer to special order)

## CROP STAGING

- Field peas and faba beans: 3–6 node stage
- Dry beans: After 1st trifoliate leaf has fully expanded up to the 2nd trifoliate leaf stage
- Soybeans: Cotyledon to 4 leaf stage
- See label for forage crop information

## TANK-MIX ORDER

- UAN
- Ransack

## NOTES

- Use 10 gal./ac. water volume to get good coverage on emerged weeds
- Ground application only
- Addition of UAN is recommended for grass control (810 mL/ac.) (sold separately)