

Valtera® Cereals

Valtera® Cereals herbicide gives you the flexibility and time needed at seeding by offering long-lasting, soil-active weed control when applied in the fall or early spring. With up to 8 weeks of residual protection, Valtera Cereals targets tough weeds like volunteer canola, cleavers and kochia—so you can seed into cleaner fields and start the season strong.

REGISTERED CROPS

- Spring wheat
- Winter wheat
- Durum wheat***

WEEDS CONTROLLED

- Annual sow thistle
- Canada fleabane
- Common chickweed
- Common ragweed²
- Common waterhemp²
- Cow cockle¹
- Cleavers
- Dandelion²
- Green pigweed²
- Eastern black nightshade²
- Hairy nightshade²
- Kochia²
- Lamb's-quarters²
- Palmer amaranth²
- Redroot pigweed²
- Sheperd's-purse¹
- Smartweed¹
- Stinkweed¹
- Volunteer canola*
- Wild buckwheat
- Wild mustard¹

WEEDS SUPPRESSED

- Green foxtail²
- Hemp-nettle¹
- Narrow-leaved hawk's-beard¹
- Perennial sow thistle**¹
- Russian thistle

BENEFITS

- **EARLY SEASON CONTROL** helps eliminate broadleaf weed pressure in your wheat during critical weed-free periods
- **EXTENDED SOIL ACTIVITY** takes out multiple flushes of tough weeds for up to 8 weeks, giving you more freedom at seeding
- **RESISTANCE MANAGEMENT TOOL** that helps manage weeds on the Valtera Cereals product label that are resistant to Group 2-, 4- and glyphosate-resistant herbicides
- **RE-CROPPING FLEXIBILITY** – most unregistered crops can be seeded 12-months after application to a registered crop
- Apply in fall or spring (check product label for specific crop registrations)
- Extended early season control provides in-crop herbicide flexibility that helps to improve the efficacy of in-crop herbicide applications by targeting fewer, smaller weeds

PERFORMANCE TIPS

1. **Recommended seeding depth** following Valtera Cereals application is 1-2 inches deep
2. **Fall application** – with a simple application in the fall when soil temperatures fall to 10 °C or less, Valtera Cereals gives you significantly more flexibility and time for your spring weed management program
3. **DO NOT** incorporate after application or prior to seeding

For specific crop usage information see label for details.

* Including herbicide-tolerant canola varieties except Clearfield® canola.

** Applications made at advanced leaf stages will reduce product effectiveness.

*** Must be applied 30 days prior to seeding.

¹ emerged weeds only (2-4 leaf)

² pre-emerge weeds only

ACTIVE INGREDIENTS

- Group 2 (florasulam)
- Group 14 (flumioxazin)

RAINFEST

- Rain required for soil activation
- For burndown efficacy, avoid application if heavy rainfall is imminent

TREATMENT AREA

- One case (2 x 8 L) treats 160 acres

APPLICATION TIMING

- For fall application, apply as late as possible but prior to freeze up for any registered crop. Ideally when the soil temperature has dropped to below 10 °C.
- Applications should be applied a minimum of 7 days prior to seeding wheat (spring, winter)
- Must be applied 30 days prior to seeding durum wheat

TANK-MIX ORDER

- Valtera Cereals
- BlackHawk® EVO (if used)
- Glyphosate

NOTES

- Seeding after application requires the use of minimum to zero-till drills
- Valtera Cereals herbicide requires moisture to activate in the soil



1.800.868.5444 | nufarm.ca

Always read and follow label directions.

BlackHawk® is a registered trademark of Nufarm Agriculture Inc.

Valtera® is a registered trademark of Valent U.S.A. LLC.

Nufarm Agriculture Inc. is a distributor of Valent Canada Inc. products in Canada.

