



A pre-emergent soil active herbicide that offers extended broadleaf weed control ahead of wheat and pulses.

REGISTERED CROPS

- Wheat (spring and durum)
- Field peas
- Soybeans
- Chickpeas
- Flax
- Sunflower

WEEDS CONTROLLED

- Common groundsel
- Common purslane
- Common waterhemp
- Eastern black nightshade
- Kochia
- Lamb's-quarters
- Large crabgrass
- Powell pigweed
- Redroot pigweed
- Russian thistle (suppression)
- Smooth crabgrass
- Wild buckwheat
- Yellow woodsorrel

BENEFITS

- **EARLY SEASON CONTROL** helps eliminate broadleaf weed pressure in pulses, wheat and soybeans during critical growth periods
- **EXTENDED SOIL ACTIVITY** for residual control of flushing weeds
- **RESISTANCE MANAGEMENT TOOL** that helps manage weeds on the Astir® product label that are Group 2-, 4- and glyphosate-resistant
- **FLEXIBLE RATE** to fit your soil type and crop program
- **FAST ABILITY TO SEED** with pre-seed or pre-emergence application timings

For specific crop usage information see label for details.

ACTIVE INGREDIENT

- Group 14 (sulfentrazone)

RAINFAST

- Avoid application if heavy rainfall is imminent

TREATMENT AREA

- One case (2 x 3.52 L) treats 80 acres at 88 mL/ac. or 60 acres at 118 mL/ac.

APPLICATION

- Tank-mix with glyphosate (if used) and/or BlackHawk® EVO/ GoldWing® (if used) for enhanced control of emerged weeds

APPLICATION TIMING

- Only apply at 88 mL/ac. rate ahead of wheat
- Fall applications can be made until freeze up
- In the spring, apply a maximum of 3 days after seeding, prior to germination

TANK-MIX ORDER

- Astir
- BlackHawk EVO or GoldWing (if used)
- Glyphosate

NOTES

- Read label for re-cropping restrictions
- For best results, use higher water volumes for good coverage
- Only use Astir or other products containing sulfentrazone once in a 24 month period



1.800.868.5444 | nufarm.ca

Always read and follow label directions.

Astir®, BlackHawk® and GoldWing® are registered trademarks of Nufarm Agriculture Inc.