

# Fierce® EZ

A pre-emergent soil active herbicide that offers extended activity on emerging spring broadleaf and grassy weeds for up to 8 weeks after activation.

## REGISTERED CROPS

- Chickpeas
- Field peas
- Lentils (fall application prior to large green and small red varieties)
- Soybeans
- Spring wheat (not including durum)
- Winter wheat

## WEEDS CONTROLLED

- Annual sow thistle
- Barnyard grass
- Chickweed
- Cleavers
- Dandelion
- Downy brome (suppression)
- Green foxtail (including Group 1- and 2-resistant biotypes)
- Green and redroot pigweed
- Japanese brome (suppression)
- Kochia (including Group 2-, 4- and glyphosate-resistant biotypes)
- Lamb's-quarters
- Russian thistle (suppression)
- Wild mustard (including Group 2-resistant biotypes)
- Wild oats (suppression)
- Volunteer canola (suppression)

## BENEFITS

- **BETTER PERFORMANCE** than other leading soil active herbicides on tough broadleaf weeds
- **ACTIVITY ON GRASSY WEEDS** with a different mode of action to manage Group 1- and 2-resistant biotypes
- **EXTENDED CONTROL** with activity that lasts up to 8 weeks
- **TWO ACTIVE INGREDIENTS** (Group 14 and 15) for management of Group 1-, 2-, 4-, 5- and glyphosate-resistant weeds
- Convenient application eliminates the need for granular broadcast herbicide products

## PERFORMANCE TIPS

1. **Emerged weeds** – use with a burndown herbicide partner like glyphosate and BlackHawk® EVO or GoldWing®
2. **Application flexibility** – Fierce® EZ alone can be applied up to 3 days post seeding soybeans, chickpeas and field peas
3. **DO NOT** incorporate after application or prior to seeding
4. Pre-seeding applications require the use of minimum disturbance seeding systems
5. Leave a small checkstrip in the field to show performance

**For specific crop usage information see label for details.**

## ACTIVE INGREDIENTS

- Group 14 (flumioxazin)
- Group 15 (pyroxasulfone)

## TREATMENT AREA

- One case (2 x 7.16 L) treats 80 acres at 180 mL/ac.

## APPLICATION TIMING

- Fall apply as late as possible but prior to freeze up for any registered crop
- Pre-emergent for soybeans, chickpeas and field peas
- Should be applied a minimum of 7 days prior to seeding spring wheat and winter wheat
- Apply up to 3 days after seeding soybeans, chickpeas and field peas

## TANK-MIX ORDER

- Fierce EZ
- BlackHawk EVO or GoldWing (if used)
- Glyphosate

## NOTES

- Seeding after application requires the use of minimum to zero till drills
- Fierce EZ requires moisture to activate in the soil



1.800.868.5444 | nufarm.ca

Always read and follow label directions.

BlackHawk® and GoldWing® are registered trademarks of Nufarm Agriculture Inc. Fierce® is a registered trademark of Valent U.S.A. LLC.

Nufarm Agriculture Inc. is a distributor of Valent Canada Inc. products in Canada.

