

Bifecta® EZ

Bifecta® EZ soil active herbicide delivers powerful, convenient pre-emergent multi-mode of action broadleaf weed control to help your crop get off to a strong start with up to 8 weeks of residual activity.

REGISTERED CROPS (FALL APPLICATION ONLY)

- Field peas
- Soybeans
- Lentils
- Spring wheat

WEEDS CONTROLLED

- Barnyard grass
- Kochia (including Group 2-, 4-, and glyphosate- resistant biotypes)
- Canada fleabane
- Lamb's-quarters
- Chickweed
- Palmer amaranth
- Common ragweed
- Russian thistle
- Dandelion
- Shepherd's-purse
- Green and redroot pigweed (including Group 2- and 5-resistant biotypes)
- Waterhemp
- Green and yellow foxtail
- Wild buckwheat
- Hairy nightshade
- Wild mustard
- Yellow woodsorrel

WEEDS SUPPRESSED

- Cleavers
- Volunteer canola (suppression including all herbicide-tolerant biotypes)

BENEFITS

- **PROVEN RESISTANCE MANAGEMENT** with multiple modes of action that target broadleaf weeds
- **EXTENDED CONTROL** with activity that lasts up to 8 weeks
- **TWO ACTIVE INGREDIENTS** (Group 5 and 14) for management of resistant weeds

PERFORMANCE TIPS

1. **Fall application** – apply in the fall when soil temperatures fall to 10 °C or less, Bifecta EZ gives you significantly more flexibility and time for your spring weed management program
2. Recommended seeding depth following Bifecta EZ application is 1-2 inches deep
3. **DO NOT** incorporate after application or prior to seeding
4. Leave a small checkstrip in the field to show performance

For specific crop usage information see label for details.

ACTIVE INGREDIENT

- Group 5 (metribuzin)
- Group 14 (flumioxazin)

TREATMENT AREA

- One case (2 x 7.13 L) treats 40 acres at 357 mL/ac.

APPLICATION TIMING

- Fall apply as late as possible, but prior to freeze up for any registered crop

TANK-MIX ORDER

- Bifecta EZ
- BlackHawk® EVO (if used)
- Glyphosate

NOTES

- Bifecta EZ requires moisture to activate in the soil
- Seeding after application requires the use of minimum to zero-till drills