

Fierce® MTZ

Fierce® MTZ soil active herbicide delivers powerful, convenient pre-emergent weed control to help your crop get off to a strong start with up to 8 weeks of residual activity on the toughest of weeds including resistant kochia and wild oats.

REGISTERED CROPS (FALL APPLICATION ONLY)

- Chickpeas
- Field peas
- Lentils
- Soybeans
- Spring wheat

WEEDS CONTROLLED

- Annual sow thistle
- Barnyard grass
- Canada fleabane
- Chickweed
- Cleavers
- Common ragweed
- Corn spurry
- Dandelion
- Downy brome
- Green foxtail (including Group 1- and 2-resistant biotypes)
- Green and redroot pigweed
- Hairy nightshade
- Japanese brome
- Kochia (including Group 2-, 4-, and glyphosate-resistant biotypes)
- Lamb's-quarters
- Night flowering catchfly
- Palmer amaranth
- Russian thistle*
- Stinkweed
- Volunteer canola (including glyphosate, glufosinate ammonium and imidazolinone tolerant varieties)
- Waterhemp
- Wild buckwheat
- Wild mustard (including Group 2-resistant biotypes)
- Wild oats

WEEDS SUPPRESSED

- Foxtail barley
- Yellow foxtail

BENEFITS

- **SUPERIOR, CONSISTENT PERFORMANCE** than other soil active herbicides on the toughest broadleaf weeds
- **PROVEN RESISTANCE MANAGEMENT** with multiple modes of action that target broadleaf weeds
- **INCREASED ACTIVITY ON GRASSY WEEDS** with modes of action to manage Group 1- and 2-resistant biotypes
- **EXTENDED CONTROL** with activity that lasts up to 8 weeks
- **THREE ACTIVE INGREDIENTS** (Group 5, 14 and 15) for management of resistant weeds

PERFORMANCE TIPS

1. **Fall application** – apply in the fall when soil temperatures fall to 10 °C or less, Fierce MTZ gives you significantly more flexibility and time for your spring weed management program
2. Recommended seeding depth following a Fierce MTZ application is 1-2 inches deep
3. DO NOT incorporate after application or prior to seeding
4. Leave a small checkstrip in the field to show performance

For specific crop usage information see label for details.

ACTIVE INGREDIENT

- Group 5 (metribuzin)
- Group 14 (flumioxazin)
- Group 15 (pyroxasulfone)

TREATMENT AREA

- One case (2 x 9.46 L) treats 40 acres at 473 mL/ac.

APPLICATION TIMING

- Fall apply as late as possible, but prior to freeze up for any registered crop

TANK-MIX ORDER

- Fierce MTZ
- BlackHawk® EVO (if used)
- Glyphosate

NOTES

- Fierce MTZ requires moisture to activate in the soil
- Seeding after application requires the use of minimum to zero-till drills

*Control at high rate.



1.800.868.5444 | nufarm.ca

Always read and follow label directions.
BlackHawk® is a registered trademark of Nufarm Agriculture Inc.
Fierce® is a registered trademark of Valent U.S.A. LLC.
Nufarm Agriculture Inc. is a distributor of Valent Canada Inc. products in Canada.

