

# Latro<sup>®</sup> WG

For post-emergent control of perennial and annual grasses and broadleaf weeds in grain and silage maize crops.

Now registered for the control of yellow bristle grass.



Control of yellow bristle grass for maize crops

## ACTIVE INGREDIENT

750g/kg nicosulfuron

## FORMULATION

Water Dispersible Granule

## BENEFITS

- The only nicosulfuron formulation registered for yellow bristle grass control
- Effective at controlling couch and mercer grass
- Also controls summer grasses and barnyard grass
- An easy to measure and mix granular formulation
- Earns Priority Partnership points

## POST EMERGENCE WEED CONTROL

Good weed control is essential when growing maize. It eliminates competition for light, nutrients and moisture, enabling the maize crop to maximise its yield. Using Latro WG as part of your post emergence weed control programme allows you to reach your full potential.

## HOW LATRO WG WORKS

Latro WG is rapidly absorbed by foliage and translocated throughout the plant. Grass and broadleaf weeds stop growing immediately, discolouration occurs followed by dieback that spreads slowly throughout the plant.

## PACK SIZE AVAILABLE

500g

## DIRECTIONS FOR USE

Grass and broadleaf weeds - Use Latro WG at 50-110g per hectare (check label for correct rate for specific weeds). Add Bonza<sup>®</sup> spray adjuvant at 500mL per 100 litres of spray volume. Apply evenly to the weed canopy in 100-300 litres of water per hectare. Use the high water rate where weeds are dense.

## WEATHER CONDITIONS

Warm, moist conditions and adequate soil moisture, both before and after application, will improve control. Maize stressed due to weather conditions may show a temporary yellowing or crop stunting.

Latro WG is rainfast six hours after application.



## APPLICATION TIMING

### Grass Weeds

Apply when there is sufficient leaf area for good uptake but before the grass has formed a thick layer. For best results with summer grass, application should be at a maximum of 3 tillers or 75mm in length and ensure Bonza is used at 1L/100L. The maize crop growth stage best for application is between the two and eight leaf stage when the crop is growing vigorously without moisture stress. Application to maize greater than 500mm in height risks crop injury.

### Broadleaf Weeds

Use Latro WG before weeds have reached the 4 true leaf stage. Weeds larger than 4 true leaf stage may only be suppressed.

## COMPATIBILITY

Latro WG is compatible in tank mixtures with Nu-Trazine™ 900DF and Flowable Atrazine 500. Latro WG may be tank mixed with Kamba® 500 (applied to maize less than 30cm tall), Emblem® or Valdo® 800WG to increase broadleaf weed control, but Bonza spray adjuvant should not be used with these mixtures as there is a greater risk of crop injury (which can be minimized by the use of dropper nozzles). Grass weed control with Latro WG may be reduced when Bonza is not used.

## WIDE WEED SPECTRUM

| Weeds                  | Control | Suppress | Weeds                          | Control | Suppress |
|------------------------|---------|----------|--------------------------------|---------|----------|
| Amaranthus (Red root)  | C       |          | Kikuyu                         | C       |          |
| Apple of peru          |         | S        | Mercer grass                   |         | S        |
| Barnyard grass         | C       |          | Oxalis                         |         | S        |
| Bathhurst bur          | C       |          | Pasture grasses (volunteers)   | C       |          |
| Black nightshade       | C       |          | Potatoes (volunteers)          |         | S        |
| Bristle grass          | C       |          | Rough bristle grass            | C       |          |
| Bindweed (convulvulus) |         | S        | Scotch thistle                 |         | S        |
| Buttercup              |         | S        | Smooth witchgrass              | C       |          |
| Californian thistle    |         | S        | Sow thistle                    | C       |          |
| Chicory                |         | S        | Summer grass                   | C       |          |
| Clover                 |         | S        | Thorn apple                    | C       |          |
| Couch                  | C       |          | Twincross                      | C       |          |
| Crowfoot grass         | C       |          | Water pepper                   | C       |          |
| Dock                   |         | S        | Wild portulaca                 | C       |          |
| Fathen                 | C       |          | Wild oats                      | C       |          |
| Fumitory               | C       |          | Willow weed                    | C       |          |
| Groundsel              | C       |          | Yellow bristle grass           | C       |          |
|                        |         |          | Key: C=Controlled S=Suppressed |         |          |

The information contained is a quick reference only. Always consult the product label before use.™Nu-Trazine is a trademark of Nufarm Ltd. ©Kamba and Valdo are registered trademarks of Nufarm Ltd. ©Emblem is a registered trademark of Nufarm SA France. ©Latro is a registered trademark of Nufarm Limited. © Nufarm Ltd 2017.

6 Manu Street, Otahuhu, PO Box 22-407, Auckland 1640, New Zealand  
**t:** 09 270 4150 **e:** nzinfo@nufarm.com **w:** nufarm.co.nz



Grow a better tomorrow.