

WARNING – Combustible liquid
HARMFUL • ECOTOXIC
KEEP OUT OF REACH OF CHILDREN

Nufarm  **Conquest**®

FREE RECYCLING



AGRECOVERY

www.agrecovery.co.nz



9 414614 017537

01-17/PS/A0701-20LNZ1.2

Active Ingredient: 100g/litre picloram as the amine salt and 300g/litre triclopyr as the butoxyethyl ester in the form of an emulsifiable concentrate.

For the control of gorse, broom, blackberry and other brushweeds, and for the spot treatment of many broadleaf weeds including ragwort, thistles and docks. Also effective for release spraying in radiata pine, and broadleaf weed control in turf.

Net: 20 Litres



Grow a better tomorrow.



HAZARD CLASSIFICATION: 3.1D, 6.1D, 6.3B, 6.4A, 6.5B, 6.9B, 9.1A, 9.2A, 9.3C
APPROVED HANDLER: This product must be under the control of an approved handler when it is applied in a wide dispersive manner or used by a commercial contractor. This product must not be used for any purpose, or in any manner, contrary to this label unless authorised under appropriate legislation.

FLAMMABLE
 • Warning – combustible liquid. Keep away from sources of ignition.

TOXICITY
 • Harmful – may be harmful if swallowed, inhaled or absorbed through the skin. May cause eye irritation. May cause mild skin irritation, or sensitisation from prolonged skin contact. May cause kidney or liver damage from repeated oral exposure at high doses.

ECOTOXIC
 • Very toxic to aquatic organisms. Avoid contamination of any water supply with product or empty container. Do not use Conquest on the bottoms of any drains, ditches or irrigation channels.
 • Selective herbicide – very toxic to some plant species. Avoid use over or near desirable plants or in areas where their roots may extend or in locations where the chemical may be washed or moved into their roots.

Beware: Spray drift hazard. Apply this product carefully. Spray drift may cause serious damage to other desirable plants. Do not allow drift to occur outside the target area.

HANDLING PRECAUTIONS STORAGE
 • Do not store near heat or flame.
 • Store in original container tightly closed and in a locked, dry, cool area away from sources of ignition and foodstuffs, seeds and fertilisers. Store in accordance with NZS 8409 Management of Agrichemicals. Do not store with Class 1, 2, 3, 2, 4 or 5 substances.
 • Stores containing 500 litres of Conquest must have two fire extinguishers.
 • Stores containing 100 litres of Conquest require secondary containment and signage, and more than 100 litres require emergency response plans.

PERSONAL PROTECTION
 • Avoid contact with skin and eyes and inhalation of spray mist.
 • When mixing or applying wear eye protection, waterproof gloves and boots, and cotton overalls. Do not eat, drink or smoke while using. Remove protective clothing and wash hands and face thoroughly before meals and after work. Wash protective clothing daily after work.

EQUIPMENT
 • Apply using well maintained and accurately calibrated equipment. Flush equipment thoroughly with clean water after use. If the spray equipment which has applied Conquest is to be used for the application of other chemicals to sensitive crops, ensure the unit is completely decontaminated.

DISPOSAL
 • Triple rinse empty container and add rinsate to the spray tank. Recycle empty container. Otherwise crush and bury in a suitable landfill.
 • Dispose of product only by using according to the label, or at an appropriate landfill.

First Aid: If swallowed do NOT induce vomiting. For advice contact the National Poisons Centre (0800 POISON (0800 764 786), or a doctor immediately. Begin artificial respiration if victim is not breathing. Use mouth-to-nose rather than mouth-to-mouth. If in eyes, flush with running water for 15 minutes and call a doctor. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with plenty of clean water.

Spillage: Wear appropriate protective clothing. Extinguish all ignition sources, exclude all bystanders from the vicinity of the spill, and prevent material from entering waterways or coming into contact with Class 1, 2, 3, 2, 4 or 5 substances. Absorb spills with inert material and place in waste containers. Wash area with water and absorb with further inert material. Dispose of waste safely in a suitable landfill.

Transport: Do not carry this product on a passenger service vehicle.

Record Keeping: Records of use as described in NZS 8409 Management of Agrichemicals must be kept if 3 litres or more of Conquest is applied within 24 hours.

Mixing and Compatibility: Pour the measured quantity of Conquest into the partially filled spray tank and add the remainder of the water. Agitate thoroughly while filling and spraying. Pulse™, where recommended, should be added last when the tank is nearly full, to minimise foaming. Conquest is compatible with most other commonly used herbicides.

Animal Safety: Poisonous plants may become more palatable after spraying and stock should be kept away from the area until plants have died down.

GENERAL
 It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

Conquest is absorbed by the leaves, stem and roots of plants and translocates to the growing points where it interferes with cell division and elongation. Plants must be actively growing for maximum efficacy. Rainfall within 3 hours following treatment may reduce effectiveness.

Resistance Management: Conquest is a pyridine carboxylic acid herbicide belonging to **GROUP [OR] HERBICIDE**. Some naturally occurring individual weeds may become resistant when excessive use of the same herbicide occurs. To maximise the effectiveness of herbicides, products containing alternate modes of action should be used in spray program. Consult www.resistance.nzpsps.org/ for alternate modes of action and the details of resistance management strategies (if any) for the crops listed on the label.

Re-establishment of Closures: Residues of Conquest in the soil may prevent the re-establishment of legumes such as clovers. The length of time that residues may vary according to site conditions, climate and the application rate. It is recommended that small areas be oversewn with clover 1-2 months prior to any planned sowing time, to check for possible harmful residues. Conquest should not be used on areas where susceptible crops or legumes such as clover or lucerne are to be sown shortly after application.

DIRECTIONS FOR USE

Application:
Ground: Use a handgun spraying pressure of 1000 – 2000 kPa and ensure complete coverage of both foliage and stems.

Aerial: Apply 200-400 litres of water per hectare using a half overlap, opposite pass technique. Aerial applications may require some follow-up ground spraying.

BRUSHWEEDS

BRUSHWEED	RATE		NOTES
	HANDGUN mL/100 Litres	AERIAL litres/ha	
Blackberry	500-650	10	Apply during summer and early autumn when bushes are actively growing. Use the high rate on hard-grazed and lower growing types on the East Coast of the North Island and in Northland. Control may be variable when these species are present.
Broom	250	10	Ground: Apply during periods of active growth from Nov. to Feb. When plants are slow growing or stressed (cold, drought, dusty, or salt sprayed conditions), use the high rate and add Pulse 100mL/100 litres to reduce variability. Follow-up treatment may be required.
Gorse	250-300	10	Ground: Apply during periods of active growth from Nov. to Feb. When plants are slow growing or stressed (cold, drought, dusty, or salt sprayed conditions), use the high rate and add Pulse 100mL/100 litres to reduce variability. NOTE – when applied to control gorse, Conquest will also control a wide range of other broadleaf weeds including ragwort, fennel, nettles, inkweed and thistles. Aerial: Apply Nov. to Feb. to actively growing bushes. Add Pulse 2L/ha to improve results on gorse. Do not treat hard-grazed or scattered bushes by air. Best results are achieved when treating bushes less than 3m high, with stems less than 3cm diameter. Follow-up treatment may be required.
Lupin	100	2	
Matagouri	500	12	Treat when in full flower during Nov. to Dec. Best control is obtained when moist spring conditions favour growth. Reduced results may occur under dry conditions or where lichens impede chemical uptake.
Sweet Brier	300-500	12	Apply Nov. to Jan. Ensure thorough wetting of the crown when gun spraying. Aerial application is enhanced when bushes are less than 2m high.
Tutu	300	10	Apply to fresh regrowth after frond expansion.
Woolly nightshade	250	10	Apply during active growth. Use Pulse to improve coverage.

Knapsack: Use 6ml per litre of water.

Mistblower: Use 100ml per 10 litres. For ULV use 500ml per 10 litres, with a No. 3 or No. 4 hole.

Stump Swabbing: To control barberry, old man's beard, hawthorn, woolly nightshade, wattles, willows and other brushweeds.

Mix 1 part Conquest with 20 parts water or diesel. Ensure continuous agitation when using diesel. Use a paintbrush or knapsack sprayer to thoroughly wet the freshly cut surface and any stems down to ground level.

Stem cut / injection: To control woolly nightshade.

Treat when trees are actively growing. For stem diameters less than 15cm, make two shallow hatchet cuts on opposite sides of the trunk at waist height, or below the first branch. For stems over 15cm diameter, make four equally spaced cuts around the trunk. Use a syringe to inject 1.5ml Conquest into each cut.

PASTURE WEEDS:

Spot Application

WEEDS	Knapsack or Battery Powered Motorbike Sprayers	High Volume gun and hose*	COMMENTS
	mL/10 litres	ml/100 litres	
Ragwort and thistles (except Californian thistles)	60	200	Rosette and multicrowned plants
		250	Early flowering plants.
Fennel, goat's rue, hemlock, horehound, inkweed, nettles	60	250	Apply during active growth from full leaf to early flowering. Californian thistle – retreatment may be required.
Alligator weed, field bindweed, Californian thistle, Cape weed, docks, sorrel	60	300	

*For low growing plants eg ragwort, thistles, etc. use a handgun with a No. 4 tip at 500 kPa. For larger semi-woody type plants eg inkweed, fennel, use a handgun with a No. 4 tip at 1000-2000 kPa.

Broadcast Application:

WEEDS	L/ha	COMMENTS
Ragwort and thistles Californian thistle Cape weed	2	Apply to heavy infestations only. Best results are achieved from application prior to flowering. Retreatment of Californian thistle may be required.

IN TREATED AREAS LEGUMES SUCH AS CLOVERS AND LUCERNE WILL BE SEVERELY AFFECTED.

Carpet Type Weedwiper:

For control of ragwort and Californian thistle, dilute 1 part Conquest with 40 parts water (1:40). Ensure the wiper remains saturated while in use. Two passes in opposite directions are recommended. Californian thistle may require retreatment. Drain and wash the wiper immediately after use.

FORESTRY: Aerial release of Radiata pine

WEEDS	RATE litre/ha	NOTES
Thistles, lupin, gorse, broom, Himalayan honeysuckle, tutu, buddlejia, wattles	1-2	Only use the high rate when trees are dormant and within one year of planting. Use the low rate when trees are not dormant. Some minor tree damage may occur. Brushweeds may only be suppressed and follow-up treatment may be required.

Grass Turf: To control creeping oxalis, dandelion, hydrocotyle, Onehunga weed and other broadleaf weeds on sports grounds and recreational areas. Use 2 litres/ha. Do not use clippings from treated areas for six months after treatment.

Registered pursuant to the ACVM Act 1997, No. P7550. See www.foodsafety.govt.nz for registration conditions.

Approved pursuant to the HSN0 Act 1996, Approval Code HSR000553. See www.epa.govt.nz for approval controls.

©Conquest is a registered trademark of Nufarm Australia Limited
 *Pulse is the registered trademark of Nufarm Technologies, USA.

CONDITIONS OF SALE:
CAUTION: READ THIS NOTICE BEFORE OPENING THE CONTAINER
If you open this container you will be deemed to have accepted the conditions below.
 The results from use of this product may be affected by factors beyond Nufarm's control, including mixing, use, climatic conditions, time of application, crop or crop stage and the possible development of resistance to the active ingredient, even though the product is reasonably fit for the purposes stated in this label, and even though label directions are followed. Nufarm will replace, free of charge, any product which is found to have a manufacturing defect. To the extent permitted by law, all other warranties, guarantees, representations and conditions, whether expressed or implied by law, trade, custom or otherwise are expressly excluded.
 When you use this product, you agree that neither Nufarm Limited nor any manufacturer of any component of the product will be liable for any damage, injury or loss including consequential loss which arises in any way from the storage, handling, or use of the product. This exclusion of liability includes loss caused by negligent acts or omissions of Nufarm Limited or any supplier or manufacturer of the product or any of their employees and agents.

Marketed by: Nufarm Limited
 6 Manu Street, Otahuhu, Auckland, New Zealand
 Telephone 0-9-270 4157
www.nufarm.co.nz
 Nufarm Emergency PhoneLink: 0800-651 911 toll free 24 hours

Bath No.:
 D.O.M.:

PROPER SHIPPING NAME			
ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S. (picloram 10%, triclopyr 30%)			
HAZARD CLASS	UN NO.	HAZCHEM	PACKING GROUP
9	3082	2X	III