

# HARMFUL • ECOTOXIC

Keep out of reach of children



# Scrubcutter<sup>®</sup>

Active Ingredient: Contains 600g/litre triclopyr as the butoxyethyl ester in the form of an emulsifiable concentrate

*Grass friendly herbicide for the control of  
gorse, broom, blackberry and other brushweeds.  
Also for broadleaf weed control in turf.*



Net: 20 Litres

L82151207



# Scrubcutter®

## HAZARD CLASSIFICATION:

6.1D, 6.3B, 6.4A, 6.5B, 6.9B, 9.1A, 9.2A, 9.3C

## APPROVED HANDLER

This product must be under the control of an approved handler when it is applied in a wide dispersive manner, or used by a commercial contractor.

This product must not be used for any purpose, or in any manner, contrary to this label unless authorised under appropriate legislation.

## TOXICITY

• Harmful – may be harmful if swallowed, inhaled or absorbed through the skin. May cause eye irritation. May cause mild skin irritation, or sensitisation from prolonged skin contact. May cause kidney damage from repeated oral exposure at high doses.

## ECOTOXIC

- Very toxic to aquatic organisms. Avoid contamination of any water supply with product or empty container especially when treating brushweeds along the banks of waterways. Application to drains, ditches and irrigation channels must only be made when these waterways are dry.
- Selective herbicide – very toxic to some plant species.
- **Beware: Spray drift hazard. Apply this product carefully. Spray drift may cause serious damage to other desirable plants.** Do not allow spray drift to occur outside the treated property.

## HANDLING PRECAUTIONS

### STORAGE

- Store in original container tightly closed and in a locked, dry, cool area away from foodstuffs, seeds and fertilizers. Store in accordance with NZS 8409 Management of Agrichemicals.
- Stores containing 100 litres are subject to signage and secondary containment and more than 100 litres require emergency response plans.

### PERSONAL PROTECTION

- Avoid contact with skin or eyes and inhalation of spray mist. When handling wear appropriate eye protection, waterproof gloves, boots and overalls. Do not eat, drink or smoke while using. Remove protective clothing and wash hands and face thoroughly before meals and after work.

### EQUIPMENT

- Apply using well maintained and accurately calibrated equipment.
- Clean equipment thoroughly with clean water after use. Do not use equipment which has applied Scrubcutter for applying other materials to desirable plants unless it has been thoroughly decontaminated using procedures recommended for phenoxy herbicides.

### DISPOSAL

- Triple rinse container and add rinsate to the spray tank. Recycle empty container. Otherwise crush and bury in a suitable landfill.
- Dispose of product only by using according to the label, or at an appropriate landfill.

**First Aid:** If swallowed do NOT induce vomiting. For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water. If splashed in eyes, wash out immediately with water. If inhaled, move the victim to fresh air immediately. Begin artificial respiration if breathing has stopped.

**Spillage:** Wear appropriate protective clothing and prevent material from entering waterways. Contain liquid spills and absorb with inert material. Wash area with water and absorb with further inert material. Dispose of waste safely in a suitable landfill.

**Transport:** Do not carry this product on a passenger service vehicle.

**Record Keeping:** Records of use, as described in NZS 8409 Management of Agrichemicals, must be kept if 3 litres or more of Scrubcutter is used within 24 hours.

**Animal Safety:** Poisonous plants may become more palatable after spraying. Stock should be kept away from the area until the plants have died down.

**Mixing and Compatibility:** Partially fill the spray tank and add the correct quantity of Scrubcutter. Agitate thoroughly while filling and spraying. Scrubcutter is compatible with most commonly used herbicides. The use of Contact™ Low Foam wetting agent is recommended to improve coverage under dry or dusty conditions.

**Shelf Life:** When stored appropriately this product should show no significant degradation for 2 years from the date of manufacture. Contact your supplier for further information about the use of any product that is older than this.

**General Comments:** Scrubcutter is damaging to clovers at the rates recommended for brushweed control. To minimise clover damage, hard graze to remove as much clover leaf as possible before spraying.

For best results spray brushweeds between November and February when they are actively growing. Use the higher rate for periods of less active growth between February and May. Spraying during winter dormancy or when plants are stressed (e.g. from drought, hard grazing, water logging, previous cutting or salt damage) can give variable results. Follow-up treatment may be required for complete control.

## Directions For Use:

### Gun Spraying

Completely wet the foliage and stems from all directions. Use a spraying pressure of 1000-2000KPa.

WEED	RATE/100 LITRES	NOTES
Gorse Broom	200ml 300ml	During periods of soft active growth between November and February. When spraying outside the optimal period or when plants are not actively growing. Pulse® Penetrant at 100ml/100L is recommended to improve control on gorse.
Blackberry	300ml 600ml	During periods of active growth in summer and early autumn. For hard grazed or lower growing types found in the North Island East Coast and Northland regions. Some 'difficult-to-kill' species, especially in Northland, are not well controlled. Remove canes in the following spring by burning, slashing or crushing.
Lupin	150ml	Apply during active growth from spring until late summer.
Sweet Brier	500ml	Apply from late flowering to early leaf yellowing.
Old Man's Beard	400ml	Treat November - March before leaf senescence.
Tutu	300ml	Treat regrowth after frond expansion.
Fennel	300ml	

## Aerial Application:

Apply with a calibrated aircraft using the half overlap-opposite pass technique. Use 200-400 litres water/hectare. Use low drift nozzles which minimise the production of small droplets. The use of a drift control agent is recommended. Preferred wind conditions are a light constant breeze away from any sensitive crop. Some follow-up ground treatment may be required.

WEED	RATE LITRES/HA	NOTES
Gorse Broom	10	Apply only during active growth (post flowering). Hard grazed scattered gorse should be ground treated. Best results are obtained from treating bushes less than 2m high with less than 3cm diameter stems. Use for desiccation 3-4 months before burning. To control actively growing seedlings beyond the cotyledon stage. Pulse® Penetrant at 250ml/100L is recommended to improve control on gorse.
Lupin	3	Apply during active growth from November to January.
Tutu	8	Treat fresh regrowth after frond expansion. Follow-up ground treatment may be required.
Blackberry	10	Apply during active growth in summer to early autumn. Remove canes in the following spring by crushing, burning or slashing. There will be some regrowth requiring follow-up treatment, especially in Northland and the North Island East Coast regions.

## Stump Swabbing:

Barberry, hawthorn, woolly nightshade, wattle, willow: Mix one part Scrubcutter with 20 parts water or diesel. Using a paintbrush or knapsack liberally cover the cut surface and stem to ground level. If using diesel agitate while filling and during use.

### Knapsack:

Use 60ml Scrubcutter per 10 litres water.

### Mistblower:

Use 100ml/10 litres water. For ULV attachments use 500ml/10 litres water, with a number 3 or 4 hole.

### Forestry:

Aerial release of radiata pine:  
For gorse, broom, lupins, Himalaya honeysuckle, wattles. Use 1-2 litres/ha. Apply the lower rate when young trees are not dormant. Only use the higher rate when trees are dormant and within one year of planting. Some minor tree distortion may occur. Weed suppression only will be obtained.

### Turf Grass:

To control hydrocotyle, creeping oxalis and some other broadleaf weeds. Apply 2 litres/ha. Some burning of fine turf grasses may occur.

Lawnbow: Use 10ml per 3 litres of water per 100 square metres. Apply twice in different directions.

Do not use clippings from treated areas for compost or mulch for 6 months after treatment.

Registered pursuant to the ACVM Act 1997, No. P5317. See [www.nzfsa.govt.nz/acvm/](http://www.nzfsa.govt.nz/acvm/) for registration conditions.

Approved pursuant to the HSN0 Act 1996, Approval Code HSR000829. See [www.ermanz.govt.nz](http://www.ermanz.govt.nz) for approval controls.

## CONDITIONS OF SALE

### CAUTION: READ THIS NOTICE BEFORE OPENING THE CONTAINER

If you open this container you will be deemed to have accepted the conditions below.

The results from use of this product may be affected by factors beyond Nufarm's control, including mixing, use, climatic conditions, time of application, crop or crop stage and the possible development of resistance to the active ingredient, even though the product is reasonably fit for the purposes stated in this label, and even though label directions are followed. Nufarm will replace, free of charge, any product which is found to have a manufacturing defect. To the extent permitted by law, all other warranties, guarantees, representations and conditions, whether expressed or implied by law, trade, custom or otherwise are expressly excluded.

When you use this product, you agree that neither Nufarm Limited nor any manufacturer of any component of the product will be liable for any damage, injury or loss including consequential loss which arises in any way from the storage, handling, or use of the product. This exclusion of liability includes loss caused by negligent acts or omissions of Nufarm Limited or any supplier or manufacturer of the product or any of their employees and agents.

© Scrubcutter and Pulse are registered trademarks of Nufarm Limited.

™ Contact is a trademark of Nufarm Limited

PROPER SHIPPING NAME	HAZARD CLASS	UN No.	HAZCHEM	PACKING GROUP
<b>ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S. (triclopryr 60%)</b>	<b>9</b>	<b>3082</b>	<b>2X</b>	<b>III</b>

Batch No.:

D.O.M.:

Marketed by Nufarm Limited

6 Manu Street, Otahuhu, Auckland, NZ

Telephone 0-9-270 5147 Fax 0-9-270 4159

Nufarm Emergency Phonenumber: 0800 651 911 toll free 24 hours

[www.nufarm.co.nz](http://www.nufarm.co.nz)

**L82161207**