

Grass Weed Control	Seedlings (2-5 leaf) or actively regenerating rhizomes		Tillering plants or hardy rhizomatous grasses	
	L/ha	L/ha	mL/ha	mL/ha
Recommended use rate <i>(refer product labels for specific recommendations)</i>				
African feather grass			1450	
annual poa	0.5	1.0	290	
barley grass	0.5	1.0	×	×
barnyard grass	0.25	0.5	115	115 + Bonza
bristle grass	0.25	0.5	115	115 + Bonza
brown top	0.5	1.0	720	960
canary grass	0.25	0.5	×	×
cocksfoot	0.25	0.5	240	480
couch: cultivated	2.0	3.0	240	480
couch: sod bound	2.0 ^b	3.0 ^b	×	960
chewings fescue	0.5 ^b	1.0 ^b	not recommended	
hairgrass	0.5 ^b	1.0 ^b	not recommended	
Indian doab	2.0	3.0	720	960
kikuyu			720	960
mercer grass			720	960
Pampas grass			1450	
paspalum	2.0	3.0	720	960
perennial ryegrass	0.25 ^c	0.5 ^c	240 + Bonza *	480 + Bonza
phalaris	0.5	1.0	×	×
prairie grass	0.5	1.0	240	480
ratstail			720	960
ripgut brome	0.5	1.0	×	×
ryegrass	0.25	0.5	115	115 + Bonza [®]
smooth witchgrass	0.5	1.0	×	×
soft brome (goosegrass)	0.5	1.0	240	480
storksbill			100	×
summer grass	0.25	0.5	115	115 + Bonza
sweet grass	0.25	0.5	×	×
sweet vernal			240	480
tall fescue	0.5	1.0	×	×
volunteer cereals	0.25	0.5 ^c	115	115 + Bonza
wild oats	0.25	0.5	95	
Yorkshire fog	0.5 ^b	1.0 ^b	240	480

a - seedlings only

b - at these rates grass weeds may only be suppressed

c - turf ryegrass and tillered cereals may only be suppressed at these rates

* - Option to use 480mL/Ha without Bonza on perennial ryegrass seedlings

©Bonza is a registered trademark of Nufarm Ltd.

©Sequence is a registered trademark of Nufarm Australia Ltd.