

**HARMFUL • ECOTOXIC**  
**Keep out of reach of children**



**Nufarm**  
**Kyte**<sup>TM</sup> **700**<sup>WG</sup>  
herbicide

**FREE RECYCLING**



**AGRECOVERY**

[www.agrecovery.co.nz](http://www.agrecovery.co.nz)

Active ingredient: Contains 700g/kg  
imazethapyr in the form of a water  
dispersible granule

Selective herbicide for controlling a range  
of weeds in clover seed crops and lucerne

**Net: 1 KG**



9 414614 037917



Grow a better tomorrow.

**Spillage:** Wear appropriate protective clothing. Prevent material from entering waterways. Contain liquid spills and absorb with inert material. Sweep up solids without creating dust. Wash spill area with water and detergent. Absorb with further inert material and dispose of waste safely in an approved landfill.

**Transport:** Do not carry more than 3kg of this product on a passenger service vehicle.

**Record Keeping:** Records of use, as described in NZS 8409 Management of Agrichemicals, must be kept if 3kg or more is applied within 24 hours.

**Resistant weeds warning:** Kyte 700WG contains a **GROUP B HERBICIDE** belonging to the imidazolinone group of chemistry, with the inhibition of acetolactate synthase (ALS) mode of action. Naturally occurring resistant biotypes are known to exist and could eventually dominate the weed population if these herbicides are used repeatedly. To minimise this risk use strictly in accordance with label instructions and resistance management strategies at [www.resistance.nzpps.org/](http://www.resistance.nzpps.org/)

**Mixing and Compatibility:** Pour the measured quantity of Kyte 700WG into the partially filled spray tank and with the agitator running top up with the remainder of the water. Maintain agitation during spraying. When tank mixing this product with other recommended compatible products, first add the other product to the tank and mix thoroughly before adding this product.

Do NOT tank mix with selective post emergence grass herbicides such as Sequence<sup>®</sup>. After application of Kyte 700WG grasses must have resumed active growth before application of grass herbicides.

**Tank Cleaning:** Flush spray equipment thoroughly with clean water after use.

## GENERAL INFORMATION

Kyte 700WG is effective against a wide range of broadleaf weeds. For grass weeds a **follow-up** spray with a selective grass herbicide such as Sequence<sup>™</sup> may be required.

Kyte 700WG may be applied either pre- or post-emergence of the weeds. When product is applied pre-emergence of the weeds, susceptible weeds may emerge but growth will be retarded and the weeds will either die or remain stunted and not compete with the crop. Good soil moisture is essential for optimum activity.

When applied post-emergence of the weeds, they will either die or remain stunted and will not compete with the crop. When applied post-emergence of the weeds always add Contact<sup>™</sup> Xcel .

Do NOT apply to very wet soils if rain is imminent or to soils prone to water logging.

Do NOT apply to crops or weeds under stress caused by factors such as root or foliar disease, nutrient deficiencies, or extremes of temperature or moisture. DO NOT use Kyte 700WG pre emergence on seedling clover or lucerne. Should re-sowing of these crops be necessary DO NOT re-apply Kyte 700WG.

Do NOT apply Kyte 700WG within 2 hours of rainfall or irrigation

**Crop Safety:** Kyte 700WG may cause slight shortening of crop internodes and/or temporary yellowing. Plants soon recover and yield is unaffected. Poor growth conditions or application when clover is beginning reproductive growth may exacerbate symptoms.

Crop	Weeds Controlled	Rate/Ha	Notes
Clover seed crops	Chickweed, cleavers, hedge mustard	100-140g	Apply when the crop is at the 2 trifoliolate leaf stage or later.
	Apple of peru, annual poa, barnyard grass, black nightshade, catsear, catchfly, chamomiles, dandelion, dock, dove's foot, fathen, fennel, field madder, fumitory, groundsel, hawksbeard, henbit, mallow, mouse-ear chickweed, parsley piert, plantain, prostrate amaranthus, redroot, shepherd's purse, smooth witchgrass, sorrel, spurrey, stinging nettle, storksbill, summer grass, twin cress, willow weed, wireweed	140g/ha	Apply to actively growing seedling weeds in the cotyledon to 4 leaf stage. Weeds may not be totally controlled but populations will be significantly reduced and surviving plants will generally be severely retarded and not compete with the crop. Good crop growth will aid weed control.  Always add Contact Xcel at 200ml/100L water
Lucerne	As listed under "clover"	140g/ha	<u>Seedling lucerne:</u> Apply when the crop is at the 2 trifoliolate leaf stage or later.  <u>Established lucerne:</u> Apply following cutting or grazing.  Apply to actively growing seedling weeds in the cotyledon to 4 leaf stage.  Always add Contact Xcel at 200ml/100L water.

**Weeds not controlled:** sow thistle, speedwell, staggerweed

**Plantback:** Under conditions that do not favour the breakdown of this product eg prolonged dry conditions, carry-over residues in the soil may affect susceptible following crops. Environmental and agronomic factors make it impossible to eliminate all injury risks associated with rotational crops. The following minimum plantback intervals (months) should be observed.

Months After Application			
6	9	18	24
beans	barley	brassicas	all other crops
clover	maize	oats	
lucerne	ryegrass	sorghum	
peas		sweetcorn	
wheat			

**Withholding Period:**

**It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.**

Do not graze or cut for stock food for 42 days after application.

## HAZARD CLASSIFICATION

6.1D, 6.3B, 6.4A, 9.1A, 9.2A

## APPROVED HANDLER

This product must be under the control of an approved handler when applied in a wide dispersive manner or by a commercial contractor. This product must not be used for any purpose, or in any manner, contrary to this label unless authorised under appropriate legislation.

## TOXICITY

- Harmful - may be harmful if swallowed, inhaled or absorbed through the skin. May cause eye irritation or mild skin irritation.

## ECOTOXIC

- Very toxic to aquatic organisms. Avoid contamination of any water supply with product or empty container. Drift and runoff from treated areas may be hazardous to aquatic plants in adjacent aquatic sites. Do not contaminate water when disposing of equipment wash water.
- Selective herbicide – very toxic to some plant species. Certain plants may be killed or damaged from root uptake of this product.

**Beware: Apply this product carefully. Spray drift may cause serious damage to other desirable plants.** It is an offence to use this product in a manner that results in damage outside the treated property.

## HANDLING PRECAUTIONS

### STORAGE

- Store in original container tightly closed and in a locked, dry, cool area away from foodstuffs, seeds and fertilizers. Storage must be in accordance with NZS 8409 Management of Agrichemicals.
- Stores containing 100kg of Kyte 700WG require signage and more than 100kg require emergency response plans.

### PERSONAL PROTECTION

- Avoid skin or eye contact and inhalation of spray mist. When handling wear eye protection, impervious gloves, boots, hat and cotton overalls buttoned to the neck and wrist. Do not eat, drink or smoke while using. Remove protective clothing and wash hands and face thoroughly before meals and after work. Wash protective clothing daily after work.

### EQUIPMENT

- Apply with well-maintained and calibrated equipment. Clean out application equipment thoroughly after use.

### DISPOSAL

- Triple rinse empty container and add rinsate to spray tank. Recycle empty container.
- Dispose of product only by using according to the label, or at an appropriate landfill.

**First Aid:** If swallowed do NOT induce vomiting. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water. If splashed in eyes, flush with running water immediately. If skin or eye irritation persists, get medical attention.

Registered pursuant to the ACVM Act 1997, No. P9045  
See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions

Bonza® and Sequence® are registered trademarks and Contact™ is a trademark of Nufarm Limited  
Kyte® is registered trademark of Nufarm Australia Limited

## CONDITIONS OF SALE:

### CAUTION: READ THIS NOTICE BEFORE OPENING THE CONTAINER

**If you open this container you will be deemed to have accepted the conditions below.**

The results from use of this product may be affected by factors beyond Nufarm's control, including mixing, use, climatic conditions, time of application, crop or crop stage and the possible development of resistance to the active ingredient, even though the product is reasonably fit for the purposes stated in this label, and even though label directions are followed. Nufarm will replace, free of charge, any product which is found to have a manufacturing defect. To the extent permitted by law, all other warranties, guarantees, representations and conditions, whether expressed or implied by law, trade, custom or otherwise are expressly excluded.

When you use this product, you agree that neither Nufarm Limited nor any manufacturer of any component of the product will be liable for any damage, injury or loss including consequential loss which arises in any way from the storage, handling, or use of the product. This exclusion of liability includes loss caused by negligent acts or omissions of Nufarm Limited or any supplier or manufacturer of the product or any of their employees and agents.

**Marketed by:** Nufarm Limited  
6 Manu Street, Otahuhu  
Auckland, NZ  
Telephone 0-9-270 4157  
[www.nufarm.co.nz](http://www.nufarm.co.nz)

Nufarm emergency phonenumber: 0800-651 911 toll free 24 hours

## PROPER SHIPPING NAME

ENVIRONMENTALLY HAZARDOUS SUBSTANCE,  
SOLID, N.O.S. (imazethapyr 70%)

## HAZARD CLASS

9

## UN No.

3077

## HAZCHEM

2Z

## PACKING GROUP

III