

SprayWise® Application Guide

HERBICIDES	FORMULATION	RECOMMENDED SPRAY QUALITY	PREFERRED WATER VOL. FOR BOOM SPRAYING (L/HA)	PULSE® PENETRANT	CONTACT™ XCEL	BONZA®	AMIGO®	GENERAL COMMENTS
Agritone® 750	SL	COARSE-VERY COARSE	100-300	N	N	N	Y	
Archer® 750	SL	COARSE-VERY COARSE	up to 300	N	N	N	Y	
Associate® 600 WDG	WDG	COARSE-VERY COARSE	200-400 (aerial)	Y	N	N	Y but not with Pulse or any other organosilicone adjuvant	
Avadex® Xtra	EC	COARSE-VERY COARSE	50-200	N	N	N	N	
Baton® 800WSG	WSG	COARSE-VERY COARSE	100-300	N	N	N	Y	
Biffo®	SL	MED-COARSE	200-500	N	Y	N	Y	A contact herbicide - good coverage is essential
Charter® 750WDG	WDG	MED-COARSE	60-300	N	N			
Chateau®	WDG	COARSE	200-400	N	N	N	Y	
Conquest®	EC	COARSE	100-300	Y	N	N	N	
Crest® 520	EC	MEDIUM	200-400	N	N	Y	N	
CRUCIAL®	SL	MED-COARSE	100-200	Y but not with Amigo or other drift control adjuvants	N	N	Y but not with Pulse or any other organosilicone adjuvant	
Dictate™ 480	SL	MEDIUM	300-500	N	Only when specifically recommended	N	N	A contact herbicide - good coverage is essential
Director™ CS	CS	COARSE	150-400	N	N	N	N	
Debut®	SL	COARSE-VERY COARSE	up to 300	N	N	N	Y	
Dockstar®	SL	MEDIUM	150-200	N	N	N	N	Specific recommendation - no adjuvants
Duplosan® KV	SL	COARSE	150-300	N	N	N	Y	
Duplosan® Super	SL	COARSE	150-300	N	N	N	Y	
Emblem® Flo	SC	COARSE	min 200	N	N	N	N	Specific recommendation - no adjuvants
Flowable Atrazine	SC	MED-COARSE	200-400	N	N	Y aids foliar control	Y	Only use one adjuvant
Kamba® 750	SL	MED-COARSE	150-300	N	N	N	Y	
Kyte™ 700WG	WDG	MED-COARSE	200-300	N	Must be used	N	N	
Latro® WG	WDG	MEDIUM	100-300	N	N	Must be used	N	
Lion® Herbicide	SL	MED-COARSE	100-200	Y but not with Amigo or other drift control adjuvants	N	N	Y but not with Pulse or any other organosilicone adjuvant	
Lion® 490 DST	SL	MED-COARSE	100-200	Y but not with Amigo or other drift control adjuvants	N	N	Y but not with Pulse or any other organosilicone adjuvant	
Nail® EC	EC	MED-COARSE	100-200	N	N	Y for seed potato haulm desiccation	N	
Nail® 600EC	EC	MED-COARSE	100-200	N	N	Y for seed potato haulm desiccation	N	
Nufarm MCPB 400	SL	COARSE-VERY COARSE	100-300	N	N	N	Y	
Nu-Trazine™ 900DF	WDG	MED-COARSE	200-400	N	N	Y aids foliar control	Y	Only use one adjuvant
Prestige®	SL	COARSE	100-300	Y when spraying brushweeds	N	Y when spraying fodder brassicas	N	
Primiera®	SC	MED-COARSE	100-300	N	N	Must be used (except with Kamba or Emblem Flo)	N	
Ramrod®	SC	MED-COARSE	220	N	N	N	N	
Relay® Super S	EC	COARSE-VERY COARSE	100-500	N	N	N	N	
Roustabout® 840	EC	COARSE	200-300	N	N	N	N	
Scrubcutter®	EC	COARSE-VERY COARSE	200-400	Y	Y	N	Y but not with Pulse or any other organosilicone surfactants	
Sempra®	WDG	MED	100-200	N	Y	Y	N	
SeQuence®	EC	FINE-MED	100-300	N	N	Must be used	N	
Sero® 750WG		MED-COARSE	min 60	Y	N	N	N	
Sprinter® 700DS	SL	COARSE-VERY COARSE	100-300	N	N	N	Y	
Tandus® XL	EC	MEDIUM	100-200	Y	N	N	N	
Thistrol® Plus	SL	COARSE-VERY COARSE	100-300	N	N	N	Y	
Tribal™ Gold	SL	COARSE-VERY COARSE	100-300	N	N	Must be used	Y	
Trimec®	SL	MED-COARSE	200	N	N	N	N	
Valdo® 800WG	WDG	COARSE	100-300	N	N	Must be used	N	
WeedMaster® Dry	WSG	MED-COARSE	100-200	Y but not with Amigo or other drift control adjuvants	N	N	Y but not with Pulse or any other organosilicone surfactants	
WeedMaster® G360	SL	MED-COARSE	100-200	Y but not with Amigo or other drift control adjuvants	N	N	Y but not with Pulse or any other organosilicone surfactants	
WeedMaster® TS470	SL	MED-COARSE	100-200	Y but not with Amigo or other drift control adjuvants	N	N	Y but not with Pulse or any other organosilicone surfactants	
WeedMaster® TS540	SL	MED-COARSE	100-200	Y but not with Amigo or other drift control adjuvants	N	N	Y but not with Pulse or any other organosilicone surfactants	

INSECTICIDES	FORMULATION	RECOMMENDED SPRAY QUALITY	PREFERRED WATER VOL. FOR BOOM SPRAYING (L/HA)	CONTACT™ XCEL	FLUME®	AMIGO®	GENERAL COMMENTS
Attack®	EC	MEDIUM	100-300 (forage brassicas) 500-700 (vegetable brassicas)	Y	Y	UNKNOWN	When using Flume with Attack on fodder crops consider crop density when adjusting water rates. Do not add Flume to Attack when spraying fodder beet
Chlorpyrifos 500EC	EC	MEDIUM	100-200	Y	Y	UNKNOWN	Do not add Flume to Chlorpyrifos 500EC when spraying fodder beet
Dew™ 600	EW	MEDIUM	150-250	Y	N	UNKNOWN	
Dimilin® 2L	SC	FINE-MED	100-200	N	N	N	
Endgame™	WDG	MEDIUM	300-500	Y	N	N	
Grandevo® WDG	WDG	MEDIUM	N/A	N	Y	N	
Kaiso® 50WG	WDG	MEDIUM	200-1000	Y	N	N	
Nuprid® 350SC	SC	MEDIUM	up to 600L	Y	N	N	
Sheriff® 100	EC	MEDIUM	400-800	Y	N	N	

FUNGICIDES	FORMULATION	RECOMMENDED SPRAY QUALITY	PREFERRED WATER VOL. FOR BOOM SPRAYING (L/HA)	FLUME®	CONTACT™ XCEL
Barrack® Betterstick	SC	MEDIUM	300-600	N	TURF AND COTULA
Captan 600 Flo	SC	MEDIUM	300-600	N	Y
Champion®++	WDG	MEDIUM	300-600	Y	Y
Champ® WG	WDG	MEDIUM	300-600	Y	Y
Crop Care Captan 900 WG	WDG	MEDIUM	300-600	Y	Y
Intuity®	SC	MEDIUM	300-600 (1000 for white rot)	N	Y
Hornet™430SC	SC	MEDIUM	300-600 (1000 for white rot)	N	Y
Nando®	SC	MEDIUM	500-600	Y	Y
Sumiscler® 500SC	SC	MEDIUM	500-1000L/ha depending on crop	N	Y
Tazer®	SC	MEDIUM	200-500	N	Y
Thiram 40F	SC	MEDIUM	300-600	Y	Y

KEY

CS	Capsule Suspension	SL	Soluble Concentrate
EC	Emulsifiable Concentrate	WDG	Water Dispersible Granule (or Dry Flowable)
EW	Oil in Water Emulsion	WSG	Water Soluble Granule
SC	Suspension Concentrate		