



PACK
SIZE

1L

5L

20L



Archer® 750

Archer 750 is a broadleaf herbicide for control of Californian thistle, yarrow, volunteer potatoes and other thistles and weeds in forage brassicas, fodder and sugar beets, cereals, maize, pasture as a mixing partner with phenoxyes and in conservation tillage.

Active ingredient

750g/L clopyralid as the dimethylamine and monomethylamine salts in the form of a soluble concentrate.

Formulation

Soluble concentrate.

Benefits

- Suitable as a tank mix partner with Baton® 800WSG, Relay® Super S and Agritone® 750 to control large or hard to kill thistles and thistles with phenoxy tolerance/resistance.
- Excellent control of yarrow and thistles in fodder beet, sugar beet, forage brassicas, cereals, pasture and maize crops.
- Grasses, cereals, fodder beet, sugar beet, maize and forage brassicas can be drilled any time after application.

Weeds controlled

Californian thistle, yarrow, volunteer potatoes, clover, ragwort, thistles, bathurst bur, daisy, dandelion, hawksbeard, hawkbit, Onehunga weed, oxeye daisy, narrow-leaved plantain.

GROUP	4	HERBICIDE
-------	---	-----------

Compatibility

Archer 750 is compatible with most commonly used herbicides, cereal fungicides and insecticides.

Refer to product labels for applicable timing and withholding periods.

Plant back periods

Grasses, cereals, fodder beet, sugar beet, maize and forage brassicas may be drilled any time after application of Archer 750.

Do not plant susceptible crops (clover, peas, beans, potatoes, tomatoes, carrots, cucurbits) within 12 weeks of applying up to 0.2L/ha of Archer 750.

Where more than 0.2L/ha is applied do not plant susceptible crops within 6–12 months of treatment without sowing a small area first to check soil residues.



Application timing and use rates

CROP	WEEDS	RATE ml/ha	NOTES
Pasture	Difficult to control thistles	40–80	Add to Relay Super S or Agritone 750 to enhance efficacy.
Cereals	Californian thistle, yarrow	120–400	Apply after the two true leaf stage of crop and before the second node stage.
	Volunteer potatoes	400	Apply after full potato emergence but before tuber formation.
Forage brassicas	Californian thistle, yarrow	200–400	Apply after the two true leaf stage of the crop and before canopy closure.
Fodder and sugar beets	Californian thistle, yarrow	200–400	Apply after the two true leaf stage of the crop and before bulbous root formation.
Maize and sweetcorn	Californian thistle, bathurst bur, yarrow	200–400	Apply after the three expanded leaf stage of the crop and before canopy closure.
	Volunteer potatoes	400	Apply after full potato emergence but before tuber formation.
Conservation tillage	Clovers, narrow-leaved plantain, thistles	200–280	For best results apply to actively growing weeds.
	Californian thistle	400	Apply from rosette to early flower. Re-treatment may be required.
	Yarrow	400	Apply to actively growing yarrow, autumn to spring. For best results surface cultivate no less than one week after treatment.
Forestry (radiata pine)	Tree release from gorse broom and wattles	4000	Add 1000ml/ha Scrubcutter® and Pulse® Penetrant. Best control when weeds are actively growing and less than 1m tall.

No grazing withholding periods when used as directed.

For control of broadleaf weeds in other crops please refer to label.

WARNING: Do not use straw from Archer 750 treated crops for compost or mulch.

The information contained is a quick reference only.

Always consult the product label before use.



© Archer, Agritone and Baton are registered trademarks of Nufarm Australia Limited.

© Pulse is a registered trademark of Nufarm Technologies USA Pty Ltd.

© Relay and Scrubcutter are registered trademarks of Nufarm Limited.

© Nufarm NZ 2025.