



PACK
SIZE

5L

20L

Omnia™

Omnia is a selective systemic herbicide providing precision control of tough broadleaf weeds across pasture, forage brassicas and plantain systems in New Zealand. It delivers robust control of key perennial broadleaf weeds while maintaining flexibility across crop rotations.

Active ingredient

30g/L aminopyralid as the triisopropanolamine salt.

Formulation

Soluble concentrate (SL).

Use rate

Typically 1.0–2.0L/ha depending on weed species, size and use pattern.

Key benefits

- Excellent control of tough perennial broadleaf weeds including thistles, ragwort and docks.
- Highly systemic activity for improved uptake and control.
- Flexible use across pasture, brassicas, plantain and sprayout situations.
- Supports rotation flexibility.
- Strong fit within the Nufarm portfolio.

Mode of action

Pyridine carboxylic acid (synthetic auxin).

GROUP	4	HERBICIDE
-------	---	-----------

Use pattern

- Apply to actively growing weeds at early growth stages for best results.
- Rate selection is driven by weed species and size, not crop.
- Avoid application to stressed crops or weeds.
- Always use with Bonza® Gold (1L/ha) in forage brassicas and plantain.

Weeds controlled

- Ragwort
- Thistles
- Docks
- Nightshade
- Fathen
- Willow weed
- Spurrey

Withholding period

- **Forage brassicas & plantain:** 7 days grazing withholding
- **Pasture:** No grazing restriction (note palatability effects)
- **Meat:** 7 days
- **Milk:** 3 days

Application

Ground-based application. Apply to small actively growing weeds and ensure correct adjuvant use and spray coverage. Suitable for broadcast and spot spraying.

Stewardship and crop safety

- Temporary leaf cupping or growth suppression may occur; yield typically unaffected.
- Clover is highly sensitive – plantback intervals of 6–12 months depending on rate.
- Animal manure may contain residues that affect sensitive crops.
- Rotate modes of action to reduce resistance risk.

Key use considerations

- Spray early for best performance.
- Match rate to weed species and size.
- Always include Bonza Gold in-crop.
- Plan rotations carefully due to clover sensitivity.



Directions for use

USE	RATE/HA	DIRECTIONS FOR USE
Forage brassicas / plantain – post-emergence	1.0–2.0L/ha + Bonza Gold 1L/ha	Apply to small actively growing weeds. Match rate to weed size. Temporary crop effects may occur but yield typically unaffected.
Pasture – broadcast	2L/ha	Apply during active growth prior to flowering. Repeat may be needed for perennial weeds.
Pasture – spot spraying	60ml/10L or 200–300ml/100L	Target individual weeds such as ragwort and thistles.
Sprayout spike	0.5–1.0L/ha	Use with glyphosate programmes to improve broadleaf weed control prior to cropping.

The information contained is a quick reference only. Always consult the product label before use.



™ Omnia is a trademark of Nufarm Limited.

© Bonza is a registered trademark of Nufarm Limited.

© Nufarm NZ 2026.