



# POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Nufarm

**TERBAZINE 875 WG**®   
**HERBICIDE** 

ACTIVE CONSTITUENT: 875 g/kg TERBUTHYLAZINE

GROUP **C** HERBICIDE

For the control of weeds in Canola (triazine tolerant), Cereals (barley, oats, wheat), Cotton, Fallows, Pulses (chickpeas, faba beans, field peas, lentils, lupins) and Sorghum.

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT**



Grow a better tomorrow.

**RESTRAINTS**

**DO NOT** apply by air.

**DO NOT** apply to waterlogged soil.

**DO NOT** apply to the fields where slope exceeds 3%.

**DO NOT** apply if heavy rains or storms that are likely to cause surface runoff are forecast within two days of application.

**DO NOT** irrigate to the point of runoff for at least 2 days after application.

**DO NOT** use as a pre-emergence application in sorghum during the wet season in the Northern Irrigation areas of Western Australia.

**DO NOT** apply more than 1.2 kg/ha per crop.

**Spray Drift Restraints:**

Specific definitions for terms used in this section of the label can be found at [apvma.gov.au/spraydrift](http://apvma.gov.au/spraydrift)

**DO NOT** allow bystanders to come into contact with the spray cloud.

**DO NOT** apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. The buffer zones in the relevant buffer zone table/s below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

**DO NOT** apply unless the wind speed is between three and 20 kilometres per hour at the application site during the time of application.

**DO NOT** apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

**DO NOT apply by a boom sprayer unless the following requirements are met:**

- spray droplets not smaller than a COARSE spray droplet size category
- minimum distances between the application site and downwind sensitive areas (see 'Mandatory buffer zones' section of the following table titled 'Buffer zones for boom sprayers') are observed.

**Buffer zones for boom sprayers**

Application rate	Mandatory downwind buffer zones	
	Natural aquatic areas	Vegetation areas
Up to maximum label rate	60 m	30 m

**Section 1. Canola (triazine tolerant), Cereals and Pulses**

CROP	SITUATION	WEEDS	RATE (kg/ha)	CRITICAL COMMENTS
Lentils (not irrigated)	Pre-Sowing (IBS only)	Burr medic ( <i>Medicago polymorpha</i> ) Black bindweed ( <i>Fallopia convolvulus</i> ) Corn gromwell/White iron weed/ Sheep weed ( <i>Buglossoides arvensis</i> ) Dead nettle ( <i>Lamium amplexicaule</i> ) Indian hedge mustard ( <i>Sisymbrium orientale</i> ) Longhead poppy ( <i>Papaver dubium</i> ) Mint weed ( <i>Chenopodium pumilio</i> ) Ox tongue ( <i>Helminthotheca echioides</i> ) Prickly lettuce ( <i>Lactuca serriola</i> ) Shepherd's purse ( <i>Capsella bursa-pastoris</i> ) Sow thistle/Milk thistle ( <i>Sonchus oleraceus</i> ) Toadrush ( <i>Juncus bufonius</i> ) Turnip weed ( <i>Rapistrum rugosum</i> ) Wild gooseberry ( <i>Physalis angulata</i> )	0.86 to 1.20	Apply pre-sowing and Incorporate By Sowing (IBS) using knife points and press-wheels only. Ensure the seed is covered with at least 3cm and preferably 5cm of soil. Refer to the APPLICATION section of the label. Do not use on light soils (ie sandy loams to loamy sands, less than 40% clay). Use the lower rate on light weed infestations. <b>DO NOT</b> use rates higher than 0.86 kg/ha on soils with pH 8.0 and above as unacceptable crop damage may occur. The soil should be free of excessive clods, trash and deep furrows. Sufficient rainfall (20 to 30 mm) to wet the soil through the weed root zone is necessary within 2 to 3 weeks of application. NOTE: Some early crop phytotoxicity may be observed at the high rate. Heavy, intense rainfall following application may cause crop damage. Avoid over-lapping sprays and spraying out corners.
Chickpeas, Faba beans, Field peas, Lupins (not irrigated)	Pre-Sowing	Wild turnip ( <i>Brassica tournefortii</i> )	0.86 to 1.20	Use the lower rate on light soils (sandy loams to loamy sands) and the higher rate on heavier soils (loams, silt plus clay 40-60%). The soil should be free of excessive clods, trash and deep furrows. Sufficient rainfall (20 to 30 mm) to wet the soil through the weed root zone is necessary within 2 to 3 weeks of application. In Pulse crops <b>DO NOT</b> use rates higher than 0.86 kg/ha on soils with pH 8.0 and above as unacceptable crop damage may occur. NOTE: Some early crop phytotoxicity may be observed particularly on light soils. Heavy, intense rainfall following application may cause crop damage. At the higher rates, avoid over-lapping sprays and spraying-out corners.
	Post Sowing, Pre Emergence (PSPE). Apply to the soil within 2 days after sowing.	Wireweed/Hogweed ( <i>Polygonum aviculare</i> ) <b>Suppression of:</b> Annual ryegrass ( <i>Lolium rigidum</i> ) Doublegee/Spiny emex ( <i>Emex australis</i> ) Phalaris ( <i>Phalaris</i> species)	0.60 to 0.86	
Canola (TT or Triazine Tolerant varieties only)	Pre-Sowing	Silver grass ( <i>Vulpia</i> species)	0.86 to 1.20	Apply when weeds are actively growing. Apply up to the 6 leaf stage of canola. Use the higher rate on weeds 6-8 true leaves at time of application. Always add Banjo® at 500 mL/100L water. <b>DO NOT</b> add any other herbicide adjuvant or wetter.
	Post Sowing, Pre Emergence (PSPE). Apply to the soil within 2 days after sowing.	Wild oats ( <i>Avena</i> species) Wild radish ( <i>Raphanus raphanistrum</i> )	0.86 to 1.20	
	Early post Emergence		0.60 to 1.20	
Cereals (wheat, oats, barley)	Pre-sowing (IBS only)		0.86 - 1.20	Apply pre-sowing and Incorporate By Sowing (IBS) using knife points and press-wheels only. Ensure the seed is covered with at least 3 cm and preferably 5 cm of soil. Refer to the APPLICATION section of the label. Use the lower rate on light soils (sandy loams to loamy sands) and the higher rate on heavier soils (loams, silt plus clay 40-60%). <b>DO NOT</b> use rates high than 0.86 kg/ha on soils with pH 8.0 and above as unacceptable crop damage may occur. The soil should be free of excessive clods, trash and deep furrows. Sufficient rainfall (20 to 30 mm) to wet the soil through the weed zone is necessary within 2 to 3 weeks of application. NOTE: some early crop phytotoxicity may be observed on light soils. Heavy, intense rainfall may cause crop damage. At the higher rates, avoid over-lapping sprays and spraying-out corners.
Wheat and barley only		All Cereal weed claims above plus control of: Annual ryegrass ( <i>Lolium rigidum</i> )	0.86 - 1.20 + 2 L/ha TriflurX®	
		All Cereal weed claims above plus control of: Wild oats ( <i>Avena</i> species)	0.86 - 1.20 + 2 L/ha Avadex® Xtra	

## Section 2. Fallows (NSW and QLD only), Cotton and Sorghum

CROP	SITUATION	WEEDS	RATE (kg/ha)	CRITICAL COMMENTS
Cotton (irrigated or dryland Roundup Ready* or Roundup Ready Flex*)	Post-Sowing Pre-Emergence	<b>Pre-emergent control:</b> Bladder ketmia ( <i>Hibiscus trionum</i> ) Boggabri weed ( <i>Amaranthus mitchellii</i> ) Caltrop/yellow vine ( <i>Tribulus terrestris</i> ) Dwarf amaranth ( <i>Amaranthus macrocarpus</i> ) Sow thistle/Milk thistle ( <i>Sonchus oleraceus</i> ) Pigweed ( <i>Portulaca oleracea</i> ) Flax leaf fleabane ( <i>Conyza bonariensis</i> )	0.86 - 1.20	Use the higher rate on heavy infestations and for longer control. Apply after sowing, before weed or crop germination, as an overall or band treatment. The minimum band width should be 40cm. Sufficient rainfall (20 to 30 mm) to wet the soil through the weed root zone is necessary within 2 to 3 weeks of application. Heavy, intense rainfall following application may cause crop damage. <b>DO NOT</b> apply using 'open cab' ground boom application equipment. Must use closed cab ground boom application equipment.
	Directed (lay-by) application	<b>Pre-emergent control:</b> Bladder ketmia ( <i>Hibiscus trionum</i> ) Boggabri weed ( <i>Amaranthus mitchellii</i> ) Caltrop/yellow vine ( <i>Tribulus terrestris</i> ) Dwarf amaranth ( <i>Amaranthus macrocarpus</i> ) Sow thistle/Milk thistle ( <i>Sonchus oleraceus</i> ) Pigweed ( <i>Portulaca oleracea</i> ) Flax leaf fleabane ( <i>Conyza bonariensis</i> )  <b>Post-emergent control:</b> Boggabri weed ( <i>Amaranthus mitchellii</i> ) Noogoora burr ( <i>Xanthium strumarium</i> ) Sow thistle/Milk thistle ( <i>Sonchus oleraceus</i> ) Turnip weed ( <i>Rapistrum rugosum</i> ) Wild gooseberry ( <i>Physalis angulata</i> )		Apply as a directed spray to cover the area beneath cotton plants and between rows when the cotton is 30-50 cm high (typically, 16-22 nodes). Minimise, spray contact or drift onto cotton plants as injury may result. Apply before new weed germinations occur or to actively growing weeds less than 8 cm wide/high. Always add Banjo® at 500 mL/100L water for post-emergent weed control. For a wider range of emerged weeds add Nufarm CRUCIAL® Advanced Technology Herbicide or Nufarm weedmaster® DST® Dual Salt Technology Herbicide. Refer to the label for complete directions and rate selection. <b>DO NOT</b> apply after the 22 node growth stage. <b>DO NOT</b> apply using 'open cab' ground boom application equipment. Must use closed cab ground boom application equipment.
Sorghum	Pre-Sowing AND Post-Sowing Pre-Emergence (PSPE)	Bladder ketmia ( <i>Hibiscus trionum</i> ) Dwarf amaranth ( <i>Amaranthus macrocarpus</i> ) Sow thistle/Milk thistle ( <i>Sonchus oleraceus</i> ) Pigweed ( <i>Portulaca oleracea</i> )	1.20	Apply to soil pre-sowing or post-sowing. Refer to the APPLICATION section below.
	Post-Emergence	Boggabri weed ( <i>Amaranthus mitchellii</i> ) Noogoora burr ( <i>Xanthium strumarium</i> ) Sow thistle/Milk thistle ( <i>Sonchus oleraceus</i> ) Turnip weed ( <i>Rapistrum rugosum</i> ) Wild gooseberry ( <i>Physalis angulata</i> )  All weeds above plus: Annual ground cherry ( <i>Physalis angulate</i> ) Apple of Peru ( <i>Nicandra physalodes</i> ) Bathurst burr ( <i>Xanthium spinosum</i> ) Bladder ketmia ( <i>Hibiscus trionum</i> ) Cow vine/Peach vine ( <i>Ipomoea lonchophylla</i> ) Dwarf amaranth ( <i>Amaranthus macrocarpus</i> ) Giant (black) pigweed ( <i>Trianthema portulacastrum</i> ) Pigweed ( <i>Portulaca oleracea</i> ) Thorn apples ( <i>Datura spp</i> )	1.20  0.95 + 250-375 mL/ha Comet® 400	Always add Banjo® at 500 mL/100L water. Apply up to the 5-7 leaf stage of sorghum. Refer to the Comet label for complete directions and rate selection.
Application to fallows prior to planting mung beans, soybeans, barley, oats and wheat as directed in the Fallows section of the label (NSW and QLD only)	Pre-emergence	African turnip weed ( <i>Sisymbrium thellungii</i> ) Burr medic ( <i>Medicago polymorpha</i> ) Corn gromwell/White iron weed/ Sheep weed ( <i>Buglossoides arvensis</i> ) Dead nettle ( <i>Lamium amplexicaule</i> ) Indian hedge mustard ( <i>Sisymbrium orientale</i> ) Flax leaf fleabane ( <i>Conyza bonariensis</i> ) Longhead poppy ( <i>Papaver dubium</i> ) Mint weed ( <i>Chenopodium pumilio</i> ) Ox tongue ( <i>Helminthotheca echioides</i> ) Prickly lettuce ( <i>Lactuca serriola</i> ) Shepherd's purse ( <i>Capsella bursa-pastoris</i> ) Sow thistle/Milk thistle ( <i>Sonchus oleraceus</i> ) Toadrush ( <i>Juncus bufonius</i> ) Wild gooseberry ( <i>Physalis angulata</i> ) Wild radish ( <i>Raphanus raphanistrum</i> ) Wild turnip ( <i>Brassica tournefortii</i> ) Wireweed/ Hogweed ( <i>Polygonum aviculare</i> ).  <b>Suppression of:</b> Annual ryegrass ( <i>Lolium rigidum</i> ) Bladder ketmia ( <i>Hibiscus trionum</i> ) Boggabri weed ( <i>Amaranthus mitchellii</i> ) Caltrop/yellow vine ( <i>Tribulus terrestris</i> ) Doublegee/Spiny emex ( <i>Emex australis</i> ) Dwarf amaranth ( <i>Amaranthus macrocarpus</i> ) Phalaris ( <i>Phalaris species</i> ) Pigweed ( <i>Portulaca oleracea</i> ) Wild oats ( <i>Avena species</i> ) Wild radish ( <i>Raphanus raphanistrum</i> )	0.86 – 1.2	Apply to moist soil or when rain is imminent and prior to germination of weeds. Use the higher rate where the soil is covered by more than 50% trash/stubble/weeds, for heavy weed pressure or for longer weed control. If weeds are present, cultivate or apply a knockdown-herbicide prior to applying Nufarm Terbazine® 875 WG. Nufarm Terbazine® 875 WG may also be tank-mixed with certain knock-down herbicides. See Compatibility section of the label. Avoid spray overlap and double spraying on headlands and corners. Refer to the Fallows section of the label for plant-back instructions.

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIODS:****HARVEST****All crops: NOT REQUIRED WHEN USED AS DIRECTED****GRAZING****Canola, Fallows, Pulses (Chickpeas, Faba beans, Field peas, Lentils, Lupins): DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 6 WEEKS AFTER APPLICATION.****Sorghum: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER APPLICATION****Cereals (wheat, barley, oats): DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 8 WEEKS AFTER APPLICATION****Cotton: DO NOT GRAZE TREATED COTTON CROPS OR FEED STUBBLES OR TRASH TO LIVESTOCK****GENERAL INSTRUCTIONS**

Nufarm Terbazine® 875 WG can be used to selectively control weeds in the crops listed in the directions for use table and on fallows prior to establishing crops as directed in Fallows section below.

When applied pre-emergence, the product is mainly absorbed via the roots. Weeds buried by cultivation may not be effectively controlled. Heavy rainfall on light soils may cause movement of the herbicide out of the weed seed zone, resulting in reduced weed control.

**DO NOT** apply to recently burnt stubble/trash. After burning, rainfall or cultivation is required to allow Nufarm Terbazine® 875 WG to reach the soil and not be irreversibly bound by ash.

Nufarm Terbazine® 875 WG is primarily incorporated by rainfall. Sufficient rainfall (20 to 30 mm either as rainfall or irrigation) to wet the soil through the weed root zone is necessary within 2 to 3 weeks of application to make the product effective. Weeds may germinate and emerge before being controlled by Nufarm Terbazine® 875 WG, however a delay in rainfall beyond 3 weeks may result in weeds germinating from depth and surviving.

Where soil has a high potential for leaching, heavy rainfall between application and crop emergence may result crop damage from movement of herbicide into the crop seed zone.

In flood or furrow irrigation situations, complete and continued activation of the product may not occur due to a thin band of dry soil on the surface during or after irrigation. Mechanical incorporation after application, using light harrows to incorporate the product into the soil not more than 4cm deep is required to ensure the irrigation water activates the product. Always apply the product to an even un-ridged seedbed.

Dense weed populations and multiple germinations (especially Wild radish) may require both a pre-emergent and post emergent herbicide treatment (only one being Nufarm Terbazine® 875 WG) to give acceptable control.

**MIXING**

**DO NOT** mix or load within 20 m of any well, dam, intermittent or perennial stream. Partially fill the spray tank with water and maintain agitation while adding the required quantity of Nufarm Terbazine® 875 WG Herbicide. Agitation should continue after filling and during spraying. If spray is allowed to stand, agitate thoroughly before resuming spraying.

**APPLICATION**

Apply by ground boom only in a minimum of 50 L/ha.

**Pre-Sowing**

Nufarm Terbazine® 875 WG Herbicide can be applied to either cultivated or un-worked soil up to 7 days prior to sowing. If cultivated, the soil should be relatively level before spraying and after sowing.

**Incorporation By Sowing (IBS)**

Best results are achieved when Incorporated By Sowing, using minimum-tillage sowing equipment (fitted with knife points or blades less than 12mm wide and minimum 20cm tyne spacing) followed by press wheels. Do not use a combination of press wheels plus harrows or chains when sowing. A high level of trash (greater than 50% ground cover) may reduce weed control and hamper the sowing operation. For crop safety, maintain slow to moderate speed to avoid leaving deep furrows and avoid throwing soil into adjacent furrows. Deep furrows may also lead to reduced weed control in the crop row.

**Post Sowing Pre Emergence (PSPE)**

Apply within 2 days of sowing to soil which is relatively flat after the sowing operation. This may require trailing harrows, prickle chain or a separate rolling operation to leave the soil in an un-ridged condition after sowing and before the application.

**Post Emergence**

Nufarm Terbazine® 875 WG Herbicide can be applied to Triazine Tolerant (TT) canola up to 6-leaf stage and sorghum up to 5-7 leaf stage of the crop. Always add Banjo® at 500 mL/100L water.

**Directed (lay-by) application**

In cotton, apply as a directed spray to cover the area beneath cotton plants and between rows when the cotton is 30-50 cm high (typically, 16-22 nodes). Minimise, spray contact or drift onto cotton plants as injury may result. Apply before new weed germinations occur or to actively growing weeds less than 8 cm wide/high. Always add Banjo® at 500 mL/100L water for post-emergent weed control. For a wider range of emerged weeds add Nufarm CRUCIAL® Advanced Technology Herbicide or Nufarm weedmaster® DST® Dual Salt Technology Herbicide. Refer to the label for complete directions and rate selection.

**Fallow Application**

Nufarm Terbazine® 875 WG Herbicide can be applied to a fallow to provide pre-emergence control of certain weeds emerging prior to sowing nominated crops below. Apply to moist soil or when rain is imminent and prior to germination of weeds. If weeds are present, cultivate or apply a knockdown-herbicide prior to applying Nufarm Terbazine® 875 WG Herbicide. Nufarm Terbazine® 875 WG Herbicide may also be tank-mixed with certain knock-down herbicides. See **Compatibility** section of the label.

**Tank Mixing**

Some reduction in control may be observed on some weeds compared to the application of knock-down herbicides alone, particularly on large weeds. It is advisable that a small scale compatibility test be carried out prior to tank mixing with other products. Read and follow all label directions, restraints, plant-back periods, withholding periods, regional use restrictions and safety directions for the tank mix products. See **Compatibility** section of the label.

**FOLLOWING CROPS**

Nufarm Terbazine® 875 WG Herbicide is broken down microbial activity, which is favoured by moist soil. Soil which is intermittently dry and moist over the plant-back period may result in longer plant backs. Observe the following plant-back intervals between the date of application to a fallow or crop and the planting date of following crops thus-

Crop to be planted after application of Nufarm Terbazine® 875 WG	Nufarm Terbazine® 875 WG application rate 0.86 kg/ha	Minimum rainfall or irrigation	Nufarm Terbazine® 875 WG application rate 1.2 kg/ha	Minimum rainfall or irrigation
Chickpeas, Faba beans, Field peas, Lentils, Lupins and Canola	3 months	105 mm	4 months	140 mm
Barley, oats, wheat	4 months	140 mm	5 months	175 mm
All other crops	6 months	175 mm	6 months	175 mm
Cotton, Sorghum, TT canola	Not required			

These plant-backs are based on normal seasonal rainfall (irrigation) and need both the months and rainfall minimums. If rain (or irrigation) during the fallow or crop growing season (where used in crop) is less than 100mm over 4 months and there are isolated heavy falls, the plant-back interval may need to be longer than indicated by the rainfall totals. If there is a prolonged dry period further advice should be sought from Nufarm Australia Limited.

**CROP SAFETY**

The safety of Nufarm Terbazine® 875 WG Herbicide to crops is due to both physically placing seed below the treated soil and to biological tolerance. A crop which is slow to germinate, emerge and grow due to frost, waterlogging, insects or disease will be less tolerant of any herbicide uptake. The following factors are most important for crop safety-

- In cereals and Pulse crops ensure the seed is covered with at least 3 cm and preferably 5 cm of soil.
- In cereals and Pulse crops, do not use rates higher than 0.86 kg/ha on soils with pH 8.0 and above as unacceptable crop damage may occur.
- Heavy rain after application onto ridged soil in particular may cause crop damage from Nufarm Terbazine® 875 WG Herbicide washed into the seed furrow. At the higher rates, avoid over-lapping sprays and spraying-out corners.
- Avoid soils which are prone to water logging or where the furrows are likely to collapse after sowing.
- Avoid using fungicide seed treatments known to reduce seedling vigour, particularly if mixing Nufarm Terbazine® 875 WG Herbicide with other herbicides.
- Sowing with disc seeders may lead to un-acceptable crop damage in cereals and Pulse crops.

**RESISTANT WEEDS WARNING**

GROUP	C	HERBICIDE
-------	---	-----------

Nufarm Terbazine® 875 WG Herbicide is a member of the triazine group of herbicides. Nufarm Terbazine® 875 WG Herbicide has the inhibitor of photosynthesis at photosystem II mode of action. For weed resistance management Nufarm Terbazine® 875 WG Herbicide is a Group C Herbicide. Some naturally occurring weed biotypes resistant to Nufarm Terbazine® 875 WG Herbicide and other triazine herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Nufarm Terbazine® 875 WG Herbicide or any other Group C herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Nufarm Australia Limited accepts no liability for any losses that may result from the failure of Nufarm Terbazine® 875 WG Herbicide to control resistant weeds.

**INTEGRATED WEED MANAGEMENT STRATEGY FOR TT CANOLA**

The use of this product in TT canola is subject to the current approved "Integrated Weed Management Strategy for the use of triazine herbicides in TT canola". The Strategy encompasses: Integrated Weed Management; the specific management of crop plant volunteers and outcrossing to other plants; and triazine herbicide residue management (agronomic and environment).

Compliance with registered label directions and adoption of the principles outlined in the Strategy will assist with implementation of crop management practices that minimise the development of herbicide resistance in treated weeds; reduce the levels of triazine residues in the environment; and manage volunteer plants and outcrossing. Copies of the Strategy are available from Nufarm Australia Limited.

Where specific information and farm management assessment is required to assist with application of *Integrated Weed Management* programs on-farm, particularly in adapting strategies to local conditions, then specific expert advice should be obtained from an appropriate agronomist, consultant, chemical supplier or Departmental adviser.

**To minimise herbicide resistance:**

- Avoid dry sowing in heavily weed infested paddocks. Wait for weed germination after the opening rains in weedy paddocks. Use a pre-plant knockdown or cultivation. No weeds should be allowed to survive at this stage.
- **DO NOT** use Nufarm Terbazine® 875 WG Herbicide if the area to be treated had a triazine herbicide applied to it last season.
- Watch for escapes, especially in paddocks with a long history of Group C herbicide use.
- **DO NOT** use Group C herbicides in consecutive years.

**Resistant Weeds Reporting**

Growers should collect plant or seed samples where weeds that are normally susceptible to atrazine, simazine and terbuthylazine may be resistant, get them tested and seek professional advice.

**COMPATIBILITY**

This product is compatible with Nufarm Avadex® Xtra Herbicide, Comet® 400 Herbicide, Nufarm CRUCIAL® Advanced Technology Herbicide, Nufarm Bouncer® 960S Herbicide, BS1000, Nufarm Banjo® Spraying Oil, TriflurX®, Amicide Advance® 700, Trooper® 75-D, weedmaster® ARGO, weedmaster® DST, Shirquat® 250, Revolver®.

**RE-ENTRY PERIOD**

**DO NOT** allow entry into treated areas for 8 days. When prior entry is necessary, wear cotton overalls buttoned to the neck and wrist over normal clothing and chemical resistant gloves. Clothing must be laundered after each day's use.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

Very toxic to aquatic life. **DO NOT** contaminate wetlands or watercourses with the product or used containers. After application of Nufarm Terbazine® 875 WG Herbicide, **DO NOT** irrigate crop to the point of run-off, unless it can be retained on farm.

**TT canola: DO NOT** use or apply this product post-emergence on raised beds or where furrows have been created in the soil for the purpose or holding or channelling water. **DO NOT** use in channels or drains.

**PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS**

**DO NOT** apply or drain or flush equipment on or near desirable trees or other plants or on areas where their roots may extend or in locations where the chemical may be washed or moved into contact with their roots. **DO NOT** apply on sites where surface water from heavy rain can be expected to run off to areas containing, or to be planted with susceptible crops or plants. **DO NOT** plant crops other than those recommended on this label for at least 6 months following the most recent application of up to 1.2 kg per hectare.

#### STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Shake bag contents into spray tank until the bag is empty. **DO NOT** dispose of undiluted chemicals on site. Puncture or shred and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers or product.

#### SAFETY DIRECTIONS

May irritate the eyes. Avoid contact with eyes. When opening the container, mixing and loading and preparing spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow length chemical-resistant gloves and a half face respirator with dust cartridge or canister. When preparing the product for use by boom spray equipment, wear cotton overalls, over normal clothing, buttoned to the neck and wrist and a washable hat and elbow length chemical-resistant gloves and a half face respirator with dust cartridge or canister. When using the prepared spray wear cotton overalls buttoned to the neck and wrist (or equivalent clothing). If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use wash gloves, contaminated clothing, and respirator and if rubber, wash with detergent and warm water.

#### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

#### ADDITIONAL STATEMENTS (WHS REGULATIONS 2011)

**Harmful if swallowed.** Wash hands and exposed skin thoroughly after handling. Do not eat, drink or smoke when using this product. IF SWALLOWED: Rinse mouth.

#### SAFETY DATA SHEET

For further information refer to the Safety Data Sheet (SDS), which can be obtained from your supplier or from the Nufarm website – [www.nufarm.com.au](http://www.nufarm.com.au)

**In case of emergency: Phone 1800 033 498** Ask for shift supervisor. Toll free 24 hours.

#### CONDITIONS OF SALE

"Any provisions or rights under the Competition and Consumer Act 2010 or relevant state legislation which cannot be excluded by those statutes or by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies, expressed or implied under common law, statute or otherwise, in relation to the sale, supply, use or application of this product, are excluded. Nufarm Australia Limited and/or its affiliates ("Nufarm") shall not accept any liability whatsoever (including consequential loss), or howsoever arising (including negligence) for any damage, injury or death connected with the sale, supply, use or application of this product except for liability which cannot be excluded by statute."

Nufarm Australia Limited  
ACN 004 377 780  
103-105 Pipe Road  
Laverton North Victoria 3026  
Tel: (03) 9282 1000  
[nufarm.com.au](http://nufarm.com.au)

© All trade marks owned or used under license by Nufarm Australia Ltd.

\* Other registered trade marks.

**APVMA Approval No.: 69790/125645**