

# CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

**Nufarm**  
**tazer<sup>®</sup>xpert<sup>™</sup>**  
**FUNGICIDE**

ACTIVE CONSTITUENTS: 80g/L AZOXYSTROBIN  
31.25g/L EPOXICONAZOLE

**GROUP 11|3 FUNGICIDE**

For the control of various diseases in wheat, barley and oats  
as specified in the DIRECTIONS FOR USE TABLE.

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT.**

© <sup>™</sup> Tazer and Xpert are trademarks of Nufarm Australia Limited.



Grow a better tomorrow.

**DIRECTIONS FOR USE****RESTRAINT:**

**DO NOT** exceed a total rate of 2 L/ha of Tazer Xpert Fungicide for a growing season.

**SPRAY DRIFT RESTRAINTS**

**DO NOT** apply when wind speed is less than 3 or more than 20 kilometres per hour at the application site.

**DO NOT** apply with spray droplets smaller than a MEDIUM spray droplet size category according to nozzle manufacturer specifications that refer to ASAE S572 standard or the BCPC guideline.

**DO NOT** apply during surface temperature inversion conditions at the application site.

Users of this product **MUST make an accurate written record** of the details of each spray application within 24 hours following application and **KEEP** this record for a minimum of 2 years. The spray application details that must be recorded are: 1. Date with start and finish times of application; 2. Location address and paddock/s sprayed; 3. Full name of this product; 4. Amount of product used per hectare and number of hectares applied to; 5. Crop/situation and weed/pest; 6. Wind speed and direction during application; 7. Air temperature and humidity; 8. Nozzle brand, type, spray angle, nozzle capacity and spray system pressure measured during application; 9. Name and address of person applying this product. (Additional record details may be required by the State or territory where this product is used.).

**MANDATORY NO-SPRAY ZONES**

**DO NOT** apply if there are aquatic and wetland areas including aquacultural ponds, surface streams and rivers downwind from the application area and within the mandatory no-spray zones shown in Table 1 below.

Table 1 – No-Spray Zones for Protection of the Aquatic Environment		
FOR AERIAL APPLICATION		
Wind Speed Range at Time of Application	Downwind Mandatory No-Spray Zone	
	Fixed-Wing	Helicopter
from 3 to 8 kilometres per hour	40 meters	40 meters
from 8 to 14 kilometres per hour	60 meters	40 meters
from 14 to 20 kilometres per hour	60 meters	60 meters
FOR GROUND APPLICATION		
from 3 to 20 kilometres per hour	5 meters	

Crop	Disease Controlled	Rate	Critical Comments
Wheat	Leaf rust ( <i>Puccinia triticina</i> ) Stripe rust ( <i>Puccinia striiformis</i> ) Stem rust ( <i>Puccinia graminis</i> f.sp. <i>tritici</i> )	1-2 L/ha <b>OR</b> 500 mL/ha + Banjo® at 1% v/v	Apply when conditions favour disease development and prior to development of disease in the crop. Aim to apply between early stem elongation and ear emergence (Z32-Z59). <b>DO NOT</b> apply later than Z59.  Select higher use rates under high infection pressure or when longer residual protection is required or in situations where the use of crop oil may present compatibility concerns or risk of crop injury e.g. tank mixtures (see compatibility section). Ensure thorough coverage.
	Septoria tritici blotch ( <i>Zymoseptoria tritici</i> )	1 L/ha <b>OR</b> 500 mL/ha + Banjo® at 1% v/v	
	Septoria nodorum blotch ( <i>Stagonospora nodorum</i> ) Tan spot ( <i>Pyrenophora tritici-repentis</i> ) Powdery mildew ( <i>Blumeria graminis</i> f.sp. <i>tritici</i> )	1-2 L/ha	
Barley	Stem rust ( <i>Puccinia graminis</i> f.sp. <i>secalis</i> )	1-2 L/ha <b>OR</b> 500 mL/ha + Banjo® at 1% v/v	Apply when conditions favour disease development and prior to development of disease in the crop. Aim to apply no later than Z59. Select higher use rates under high infection pressure or when longer residual protection is required or in situations where the use of crop oil may present compatibility concerns or risk of crop injury e.g. tank mixtures (see compatibility section). Ensure thorough coverage.
	Leaf scald ( <i>Rhynchosporium secalis</i> )	1-2 L/ha	
	Net Form Net blotch ( <i>Pyrenophora teres</i> f.sp. <i>teres</i> ) Spot Form Net blotch ( <i>Pyrenophora teres</i> f.sp. <i>maculata</i> ) Leaf rust ( <i>Puccinia hordei</i> )	1-2 L/ha <b>OR</b> 500 mL – 1 L/ha + Banjo® at 1% v/v	Apply when conditions favour disease development and prior to development of disease in the crop. Aim to apply between early stem elongation and ear emergence (Z32-Z59). <b>DO NOT</b> apply later than Z59. Select higher use rates under high infection pressure or when longer residual protection is required or in situations where the use of crop oil may present compatibility concerns or risk of crop injury e.g. tank mixtures (see compatibility section). Ensure thorough coverage.
	Powdery mildew ( <i>Blumeria graminis</i> f.sp. <i>hordei</i> )		

Crop	Disease Controlled	Rate	Critical Comments
Oats	Septoria leaf blotch ( <i>Phaeosphaeria avenaria</i> )	500 mL – 1 L/ha	Apply when conditions favour development of disease but before high levels of disease develop in the canopy. Repeat sprays may be needed if infection occurs early in crop development and conditions favour infection. Regularly monitor for signs of re-infection from first application. Use the higher rates specified in the rate column under high infection pressure or when longer residual protection is required. Lower rates are effective under low disease pressure but have a reduced residual effect. Ensure thorough coverage. <b>DO NOT</b> apply later than Z59.
	Stem rust ( <i>Puccinia graminis</i> f.sp. <i>avenae</i> )	1 L/ha <b>OR</b> 500 mL/ha + Banjo® at 1% v/v	Apply when conditions favour disease development and prior to development of disease in the crop. Aim to apply no later than Z59. Select higher use rates in situations where the use of crop oil may present compatibility concerns or risk of crop injury e.g. tank mixtures (see compatibility section). Ensure thorough coverage.

## NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

### WITHHOLDING PERIODS:

**GRAZING: DO NOT GRAZE OR CUT FOR STOCKFOOD FOR 3 WEEKS AFTER APPLICATION.**

**HARVEST: NOT REQUIRED WHEN USED AS DIRECTED.**

### EXPORT OF TREATED PRODUCE

MRLs or import tolerances for azoxystrobin and epoxiconazole may not be established in all markets. If you are growing cereals for export, please check with Nufarm Australia Ltd for the latest information.

### LIVESTOCK DESTINED FOR EXPORT MARKETS

The grazing withholding period only applies to stock slaughtered for the domestic market. Some export markets apply different standards. To meet these standards, ensure that in addition to complying with the grazing withholding period, the Export Slaughter Interval is observed before stock are sold or slaughtered.

### EXPORT SLAUGHTER INTERVAL (ESI):

AFTER OBSERVING THE WITHHOLDING PERIOD FOR GRAZING OR CUTTING FOR STOCKFOOD, LIVESTOCK THAT HAVE BEEN GRAZED ON OR FED TREATED CROPS SHOULD BE PLACED ON CLEAN FEED FOR 3 DAYS PRIOR TO SLAUGHTER.

### GENERAL INSTRUCTIONS:

Nufarm Tazer Xpert Fungicide is a broad spectrum contact and systemic fungicide. It is taken up via the stem and foliage and is translocated upwards and outwards to provide protection to new growth. Thorough coverage of the crop is necessary for best results.

Check the crop regularly for any symptoms of disease infection.

### MIXING

Nufarm Tazer Xpert Fungicide is a suspension concentrate (SC) formulation. Add the required amounts of Nufarm Tazer Xpert Fungicide to the spray tank when partly filled with water and begin agitation. Ensure agitation is maintained while topping up the tank and during application.

Where the addition of Banjo is recommended ensure Nufarm Tazer Xpert Fungicide is added to the spray tank under agitation and mixed well through the spray solution prior to the addition of the Banjo. **DO NOT** allow prepared mixture to stand overnight.

### APPLICATION

#### Ground Application

Apply in a water volume of between 80 and 100 L/ha, using a MEDIUM spray quality. Use the higher water volume in crops with heavier canopies.

#### Aerial Application

Apply with suitable aircraft, set up and operated to apply fungicides to cereal crops in a minimum water volume of 20L/ha.

### FUNGICIDE RESISTANCE WARNING

GROUP	<b>11 3</b>	FUNGICIDE
-------	-------------	-----------

Nufarm Tazer Xpert Fungicide is a member of the Quinone outside Inhibitors (QoIs) and DMI groups of fungicides. For fungicide resistance management the product is a Group 11 and 3 fungicide. Some naturally occurring individual fungi resistant to the product and other Group 11 and 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product or other Group 11 and 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Nufarm Australia Limited accepts no liability for any losses that may result from the failure of this product to control resistant fungi.

### COMPATIBILITY

Tazer Xpert Fungicide is compatible with Banjo. In mixtures **without** Banjo, Tazer Xpert Fungicide is compatible with Amicide® Advance 700, Estericide® Xtra 680, Kaiso® 240EG Insecticide with Sorbie® Technology and UAN (urea ammonium nitrate such as Flexi-N\*, when UAN is 50% or less of total spray volume and when Tazer Xpert is applied by itself or in a tank mixture with Kaiso). For further information on compatibility please contact your local reseller or Nufarm Australia Limited representative.

### PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET CROPS

**DO NOT** apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

### PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Highly Toxic to Aquatic organisms. **DO NOT** contaminate streams, rivers or waterways with the chemical or used containers.

### STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Empty containers and product should **NOT** be burnt.

**SAFETY DIRECTIONS**

May irritate the eyes. Avoid contact with eyes. When opening the container and preparing the product for use and using the product, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. In addition, when preparing the product for use by aerial spraying equipment, wear cotton overalls over normal clothing buttoned to the neck and wrist and chemical resistant gloves. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use wash gloves and contaminated clothing.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26; New Zealand 0800 764 766.

**ADDITIONAL STATEMENTS (WHS REGULATIONS 2011)**

**Suspected of causing cancer. Suspected of damaging fertility or the unborn child. Causes serious eye damage.** Wear protective gloves, clothing, eye and face protection. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

**SAFETY DATA SHEET**

For further information refer to the Safety Data Sheet (SDS), which can be obtained from your supplier or from the Nufarm website – [www.nufarm.com.au](http://www.nufarm.com.au)

**In case of emergency: Phone 1800 033 498** Ask for shift supervisor. Toll free 24 hours.

**CONDITIONS OF SALE**

"Any provisions or rights under the Competition and Consumer Act 2010 or relevant State legislation which cannot be excluded by those statutes or by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies, expressed or implied under common law, statute or otherwise, in relation to the sale, supply, use or application of this product, are excluded. Nufarm Australia Limited and/or its affiliates ("Nufarm") shall not accept any liability whatsoever (including consequential loss), or howsoever arising (including negligence) for any damage, injury or death connected with the sale, supply, use or application of this product except for liability which cannot be excluded by statute."

Nufarm Australia Limited  
ACN 004 377 780  
103-105 Pipe Road,  
Laverton North Victoria 3026  
Tel: (03) 9282 1000  
[nufarm.com.au](http://nufarm.com.au)

© All trade marks owned or used under licence by Nufarm Australia Ltd.

\* Other trade marks

**APVMA Approval No.: 69803/107857**