

**CAUTION**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



ACTIVE CONSTITUENT:  
400g/kg TRALKOXYDIM

**GROUP A HERBICIDE**

For the control of wild oats and annual ryegrass and suppression of annual phalaris in wheat, barley, rye and triticale.



**Nufarm**

Grow a better tomorrow.

APVMA Approval No.: 46159/104094

**DIRECTIONS FOR USE**

**RESTRAINTS**

DO NOT apply to weeds or crops under stress due to severe frost, drought, water logging, low soil fertility, pre-emergent herbicides, insects or diseases.

DO NOT use boomless jets or misters.

DO NOT apply if rain is expected within a half-hour.

CROP	WEEDS CONTROLLED	WEED STAGE	STATE	RATE/HA	CRITICAL COMMENTS
Wheat, Barley Rye, Triticale	Wild oats	Growing actively at 2-4 leaf (pre-tillering)	Qld, NSW, Vic, Tas, SA only	300g*	Apply in first 4-6 weeks to unstressed crops and weeds when the majority of crop plants have 2 leaves. Application to lush crops may increase likelihood of yellowing and/or growth retardation.
				380g*	Apply in the first 4-6 weeks to unstressed crops and weeds when the majority of crop plants have 3 leaves. This is the minimum rate if tank mixing with listed broadleaf herbicides.
				430g*	Apply in first 4-6 weeks when the majority of crop plants have 3 leaves. Use this rate under dry soil conditions or if applying during August onwards. Use of Supercharge® at 1L per 100L may improve weed control under these conditions.
		500g*	Use where application is delayed. Do not apply to pre-tillered crops. Use of Supercharge at 1L per 100L may improve weed control if soil is dry or if applying during August onwards.		
	Wild oats and Annual ryegrass in mixed infestations	Growing actively at 2-4 leaf (pre-tillering)	NSW, Vic, Tas., SA only	380g*	Use when wild oats predominate. Do not apply to crops with less than 3 leaves. Use of Supercharge at 1L per 100L may improve weed control if soil is dry or if applying during August onwards.

CROP	WEEDS CONTROLLED	WEED STAGE	STATE	RATE/HA	CRITICAL COMMENTS
		Growing actively up to early tillering (Maximum of 2 tillers)		500g*	Do not apply to pre-tillered crops. Use of Supercharge at 1L per 100L may improve weed control of soil is dry or if applying during August onwards.
		Growing actively at 2-4 leaf (pre-tillering)		380-500g*	Do not apply to tillered ryegrass. Using spray volumes of 70-100L/ha may assist control. Tank mixes of the lower rate with listed broadleaved herbicides may reduce ryegrass control.
Wheat, Barley, Rye, Triticale	Annual phalaris (suppression only of Paradoxa grass, Lesser canary grass)	Growing actively at 2-4 leaf (pre-tillering)	NSW, Vic, Tas., SA only	380-500g*	Preferably apply within 6 weeks of sowing when weeds are at the 2 to 4 leaf stage and growing actively. Do not apply to tillered annual phalaris. The higher rate may give more reliable results. Use of spray volumes of 70-100L/ha may assist control. Tank mixes with listed broadleaved herbicides may reduce annual phalaris control. Paradoxa grass is more sensitive than lesser canary grass. Do not apply to crops with less than 3 leaves.
Wheat, Barley, (except Moondyne Yagan), Rye, Triticale	Wild oats	Growing actively at 2-4 leaf (pre-tillering)	WA only	250g*	Apply in first 4-6 weeks to unstressed crops and weeds when the majority of crop plants are at growth stage Z12. Application to lush crops may increase likelihood of yellowing and/or growth retardation.
		Growing actively up to early tillering (Maximum of 2 tillers)		380g*	Use where application is delayed and the majority of crop plants are at growth stage Z13. This is the preferred rate for dense wild oat populations. Use this rate if tank mixing with listed broadleaf herbicides.
	Wild oats and Annual ryegrass in mixed infestations	Growing actively up to early tillering (Maximum of 2 tillers)		380g*	Use when wild oats predominate. Apply when the majority of crop plants are at growth stage Z13.
		Growing actively at 2-4 leaf (pre-tillering)			Do not apply to tillered ryegrass. Use of higher spray volumes may assist control. Tank mixes with listed broadleaved herbicides may reduce ryegrass control.

**\*ALWAYS ADD SUPERCHARGE. THE NORMAL RATE IS 750mL PER 100L SPRAY SOLUTION. INCREASE THE RATE TO 1L PER 100L UNDER DRYING CONDITIONS. ALWAYS USE 1L PER 100L FOR AERIAL APPLICATION.**

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

#### WITHHOLDING PERIOD

**DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION.**

#### GENERAL INSTRUCTIONS

##### MIXING

Only use sprayers in good working condition with high capacity agitation. Ensure the sprayer is cleaned according to the label instructions on the product used prior to Achieve WG. Use only clean water for the spray solution. Ensure that all in-line strainer and nozzle screens in the sprayer are 100 mesh or coarser. Do NOT use cartridge or disc filters after the spray tank outlet, or have two (or more) filters directly after each other anywhere in the system.

Begin to fill spray tank with clean water, and engage agitator. Agitation must be continued throughout the entire mixing and spraying procedure. When the sprayer is half full of water, add Achieve WG slowly to the spray tank through the tank opening or through a hopper designed to handle granules. The hopper should be dry before adding Achieve WG granules.

- Do not premix Achieve WG.
- Add the total required amount of Achieve WG from all cartons first, prior to adding other herbicides or Supercharge adjuvant.
- Wait at least one minute after the last of the Achieve WG has been added to the tank to allow for complete dispersion of the granules.
- Add the recommended broadleaf weed herbicide or insecticide next if tank mixing.
- Slowly add Supercharge to the tank last, in at least 10 times its volume of water especially where a suction (filling) probe or hopper is used.
- Continue to fill tank to desired level with water.

#### SPRAYING INSTRUCTIONS

##### Ground equipment

Apply in a spray volume of 50-150 litres of water per hectare at a pressure of 250-350 kPa to ensure proper dispersion and spraying characteristics of the Achieve WG spray mix.

Use flat fan nozzles such as Spraying Systems 110-01, 110-015, 110-02 or equivalent sizes from other manufacturers. Do not use 110-03 or bigger nozzles with Achieve. Do not use floodjet nozzles, boomless jets or misters or controlled droplet application equipment. Nozzles may be angled forward 45° to enhance penetration of the crop and provide better coverage.

Always ensure that agitation is continued until spraying is completed even if stopped for brief periods of time. If the agitation is stopped for more than 5 minutes re-suspend the spray solution by running on full agitation until the settled particles are fully re-suspended, prior to spraying. The spray mixtures of Achieve WG and Supercharge may be left overnight without loss of efficacy provided the settled product is completely re-suspended by stirring and/or agitation prior to re-use.

#### Aerial equipment

Apply through boom or Micronair units in 20-30 litres of water per hectare. The use of the higher volume is preferred due to more reliable results. Droplet size should be 200-250 microns (VMD) and swath width should not be wider than recommended for type of plane and application unit. Weed control following aerial application may not be as good as from ground application.

#### CROP SAFETY

If Achieve WG is applied to stressed crops or treatment is followed by stress or if crop plants have less than 2 true leaves (Z12), yellowing or growth retardation may occur. Crops normally recover without loss in yield. Avoid double spraying (overlap).

#### PLANTBACK PERIODS

Do not plant sorghum, maize or rice for a period of four weeks after application of Achieve WG. Allow at least 10 days between application of Achieve WG and any other herbicide, except Ally for which the interval should be 14 days after Achieve WG. Allow at least 10 days between application of Achieve WG and any fertilizer treatment.

#### RESISTANT WEEDS WARNING

<b>GROUP</b>	<b>A</b>	<b>HERBICIDE</b>
--------------	----------	------------------

Achieve WG Herbicide is a member of the cyclohexanediones group of herbicides. Achieve has the inhibitors of acetyl coA carboxylase mode of action. For weed resistance management Achieve is a Group A herbicide. Some naturally occurring weed biotypes resistant to Achieve and other inhibitors of acetyl coA carboxylase herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Achieve or other inhibitors of acetyl coA carboxylase herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Nufarm Australia Limited accepts no liability for any losses that may result from the failure of Achieve to control resistant weeds.

#### COMPATIBILITY

For broadleaved weed control, Achieve WG may be mixed as below.

STATE	ACHIEVE WG g/ha	CO-HERBICIDE MAXIMUM L/ha	
Qld, NSW, Vic, SA only	380-500g	Quattro®, Buctril® MA	1-1.4
		Polo®, MCPA ester	0.5/1.0
		Tigrex®(mixture may cause temporary chlorosis)	0.8
		Lontrel®	0.3
		Eclipse®	7g
WA only	380g	Quattro®, Buctril® MA	1.0
		Polo®, MCPA ester	0.5*
		*Note restriction on Kulin.	
		Tigrex® (mixture may cause temporary chlorosis)	0.8
		Lontrel®	0.3
		Eclipse®	7 g

DO NOT mix Achieve WG with 2,4-D, MCPA amine, dicamba or Ally (metsulfuron-methyl). Allow at least 10 days between application of Achieve WG and herbicides containing 2,4-D, MCPA amine, dicamba or similar actives. Allow at least 14 days between application of Achieve WG and Ally (metsulfuron-methyl). Achieve WG should be applied prior to the broadleaf herbicides.

#### INSECTICIDES

Achieve WG is compatible with Axe, Mascot Duo, Imidan, Matador Zeon and Saboteur.

#### PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment which may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.



[spraywisedecisions.com.au](http://spraywisedecisions.com.au) is an online weather forecasting program and is recommended for use when planning your pesticide application.

#### PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Moderately hazardous to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used container.

#### STORAGE AND DISPOSAL (5 and 10kg)

Store in the closed, original container in a dry, cool, well ventilated area out of direct sunlight. Single rinse or shake remainder into spray tank. Do not dispose of undiluted chemicals on site. Puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

#### **STORAGE AND DISPOSAL (50 kg)**

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

#### **SAFETY DIRECTIONS**

Harmful if swallowed. Will irritate the eyes. If product in eyes, wash it out immediately with water. When preparing spray wear face shield or goggles. Wash hands after use. After each day's use, wash face shield or goggles.

#### **FIRST AID**

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131 126.

#### **ADDITIONAL STATEMENTS (WHS REGULATIONS 2011)**

**Suspected of causing cancer. Causes serious eye damage. Causes skin irritation.** Wash hands and exposed skin thoroughly after handling. Do not eat, drink or smoke when using this product. Wear protective gloves, clothing, eye and face protection. IF SWALLOWED: Rinse mouth. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. IF ON SKIN: Wash with plenty of soap and water. If skin irritation occurs: Get medical advice/attention. Take off contaminated clothing and wash before reuse.

#### **SAFETY DATA SHEET**

For further information refer to the Safety Data Sheet (SDS) which can be obtained from your supplier or from the Nufarm website – [nufarm.com.au](http://nufarm.com.au)

**In case of emergency: Phone 1800 033 498** Ask for shift supervisor. Toll free 24 hours.

#### **CONDITIONS OF SALE**

"Any provisions or rights under the Competition and Consumer Act 2010 or relevant state legislation which cannot be excluded by those statutes or by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies, expressed or implied under common law, statute or otherwise, in relation to the sale, supply, use or application of this product, are excluded. Nufarm Australia Limited and/or its affiliates ("Nufarm") shall not accept any liability whatsoever (including consequential loss), or howsoever arising (including negligence) for any damage, injury or death connected with the sale, supply, use or application of this product except for liability which cannot be excluded by statute."

Nufarm Australia Limited  
ACN 004 377 780  
103-105 Pipe Road  
Laverton North Victoria 3026  
Tel: (03) 9282 1000

© All trade marks owned or used under license by Nufarm Australia Limited.

\* Other trade marks



**Nufarm**

Grow a better tomorrow.

Nufarm Australia Limited  
ACN 004 377 780  
103-105 Pipe Road  
Laverton North Victoria 3026  
Tel: (03) 9282 1000  
nufarm.com.au

Net Contents: 5kg  
Net Contents: 10kg  
Net Contents: 50kg

APVMA Approval No.: 46159/104094

#### STORAGE AND DISPOSAL (5 and 10kg)

Store in the closed, original container in a dry, cool, well ventilated area out of direct sunlight. Single rinse or shake remainder into spray tank. Do not dispose of undiluted chemicals on site. Puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

#### STORAGE AND DISPOSAL (50 kg)

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

#### SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes. If product in eyes, wash it out immediately with water. When preparing spray wear face shield or goggles. Wash hands after use. After each day's use, wash face shield or goggles.

#### FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131 126.

#### ADDITIONAL STATEMENTS (WHS REGULATIONS 2011)

**Suspected of causing cancer. Causes serious eye damage. Causes skin irritation.** Wash hands and exposed skin thoroughly after handling. Do not eat, drink or smoke when using this product. Wear protective gloves, clothing, eye and face protection. IF SWALLOWED: Rinse mouth. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. IF ON SKIN: Wash with plenty of soap and water. If skin irritation occurs: Get medical advice/attention. Take off contaminated clothing and wash before reuse.

**SAFETY DATA SHEET**

For further information refer to the Safety Data Sheet (SDS), which can be obtained from your supplier or from the Nufarm website – [nufarm.com.au](http://nufarm.com.au)

**In case of emergency: Phone 1800 033 498** Ask for shift supervisor. Toll free 24 hours.

**CONDITIONS OF SALE**

“Any provisions or rights under the Competition and Consumer Act 2010 or relevant state legislation which cannot be excluded by those statutes or by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies, expressed or implied under common law, statute or otherwise, in relation to the sale, supply, use or application of this product, are excluded. Nufarm Australia Limited and/or its affiliates (“Nufarm”) shall not accept any liability whatsoever (including consequential loss), or howsoever arising (including negligence) for any damage, injury or death connected with the sale, supply, use or application of this product except for liability which cannot be excluded by statute.”



© All trade marks owned or used under license by Nufarm Australia Limited.

\* Other trade marks

BN / DOM / Barcode / /