

Nufarm Australia Limited  
 ACN 004 377 780  
 103-105 Pipe Road  
 Laverton North Victoria 3026  
 Tel: (03) 9282 1000  
 Fax: (03) 9282 1001  
 nufarm.com.au

Contents: 20 Litres  
 Contents: 15 Litres  
 Contents: 110 Litres  
 Contents: 200 Litres  
 Contents: 500 Litres  
 Contents: 1000 Litres

APVMA Approval No.: 62527/51805

#### STORAGE AND DISPOSAL (20L, 200L)

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemical on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, triple rinse, break, crush or puncture and deliver empty packaging for appropriate disposal at an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Empty containers and product must not be burnt.

#### (15L)

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Triple or preferably pressure rinse inner bladder before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean packaging to recycler or designated collection point. If not recycling, deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

#### Refillable Containers (500L & 1000L)

Store in the closed, original container in a cool well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of sale for refill or storage.

#### Envirodrum 110L Mini Bulk Returnable Container

Store in original sealed Envirodrum in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the Envirodrum with water or any other foreign matter. After each use of the product, please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase. The Envirodrum remains the property of Nufarm Australia Limited.

#### SAFETY DIRECTIONS

Harmful if swallowed. Attacks eyes. Will irritate the skin. When preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms, and face thoroughly with soap and water. After each day's use wash gloves, goggles and contaminated clothing.

#### **FIRST AID**

If poisoning occurs, contact a Doctor or Poisons Information Centre Phone Australia 13 11 26.

#### **ADDITIONAL STATEMENTS (WHS REGULATIONS 2011)**

**Harmful in contact with skin. Harmful if inhaled. Causes serious eye irritation.** Avoid breathing spray. Wash hands and exposed skin thoroughly after handling. Do not eat, drink or smoke when using this product. Wear protective gloves, clothing, eye and face protection.

#### **SAFETY DATA SHEET**


For further information refer to the Safety Data Sheet (SDS), which can be obtained from your supplier or from the Nufarm website – [www.nufarm.com.au](http://www.nufarm.com.au)

**In case of emergency: Phone 1800 033 498** Ask for shift supervisor. Toll free 24 hours.

#### **CONDITIONS OF SALE**

“Any provisions or rights under the Competition and Consumer Act 2010 or relevant state legislation which cannot be excluded by those statutes or by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies, expressed or implied under common law, statute or otherwise, in relation to the sale, supply, use or application of this product, are excluded. Nufarm Australia Limited and/or its affiliates (“Nufarm”) shall not accept any liability whatsoever (including consequential loss), or howsoever arising (including negligence) for any damage, injury or death connected with the sale, supply, use or application of this product except for liability which cannot be excluded by statute.”

BN / DOM / Barcode / /



APVMA Approval No.: 62527/51805

#### DIRECTIONS FOR USE

SITUATION & CROP	WEEDS	RATE	CRITICAL COMMENTS
Wheat (up to early flag leaf (Z37) only)	Refer Weed Table	440mL-1.84L/ha	Use the lower rate (440mL/ha) when crop is at 3-5 leaf stage (Z13-15). Apply higher rates from 5 leaf stage through to flag leaf just visible (Z15-37). Kulin: Use only from Z15-37 (WA only).
Oats, Barley, Triticale and Cereal Rye (up to early flag leaf (Z37) only)		440mL-1.4L/ha	Use lower rate (440mL/ha) when crop is at 3-5 leaf stage (Z13-15). Apply higher rates from 5 leaf stage through to flag leaf just visible (Z15-37). Barley: Use only from 5 leaf stage to flag leaf just visible (Z15-37).
Grass Pastures and Grass Seed Crops		440mL-1.84L/ha	Apply to established grass based pastures. Damage may occur to Legumes if present. See Legume Tolerance under Protection of Crops. NOTE: Capeweed, Paterson's curse (Salvation Jane) and Variegated thistle may be poisonous to stock after spraying.
CEREALS Wheat, barley, oats, triticale, cereal rye	Wild radish Stage of weed growth Up to the 4 leaf stage and not more than 120mm in diameter	LVE Agritone 175mL/ha plus Nugrex® 350mL/ha	<p>^Reduced efficacy (suppression only) may be achieved on wild radish larger than 8 leaf or greater than 180mm in diameter.</p> <p>DO NOT use this tank-mix if cereals are undersown with lucerne or annual medics.</p> <p><b>Crop Stage</b>  Nugrex 350mL/ha+ LVE Agritone 175mL/ha: Apply from 3 leaf to fully tillered (Z13-30).  Nugrex 500mL/ha + LVE Agritone 175mL/ha: Apply from 3 leaf to fully tillered (Z13-30).  Nugrex 500mL/ha + LVE Agritone 350mL/ha: Apply from 5 leaf stage to fully tillered (Z16-30).  Optimum results are achieved when sprayed at 3-5 leaf crop stage (generally 4-8 weeks post-sowing).  <b>WA only:</b> DO NOT apply to Barley or Kulin Wheat before the 5 leaf stage (Z15).  <b>Warning:</b> Nugrex may cause transient crop yellowing of cereals. Some varieties of oats have not been tested. (Refer to "Crop Tolerance" section of General Instructions on the Nugrex label).</p>
	Up to the 6 leaf stage and not more than 150mm in diameter	LVE Agritone 175mL/ha plus Nugrex 500mL/ha	
	Up to the 8 leaf stage and not more than 180mm in diameter <sup>^</sup>	LVE Agritone 350mL/ha plus Nugrex 500mL/ha	

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIOD:**

**HARVEST:** NOT REQUIRED WHEN USED AS DIRECTED.

**GRAZING:** DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

**GENERAL INSTRUCTIONS**

Before opening, carefully read Directions for Use, Precautionary Statements, Safety Directions and First Aid Instructions. Weeds should be sprayed while actively growing and at their most susceptible stage. As a general guide, annuals should be sprayed when young and perennials just prior to appearance of bud stage. Extremes of cold or drought are unfavourable. The best conditions are when soil is moist, weather fine and rain unlikely within 6 hours following application. DO NOT spray weeds outside the stages indicated in "Critical Comments" as damage, loss of yield or inadequate weed control may result. DO NOT spray in high winds.

**APPLICATION/MIXING INFORMATION**

This product mixes readily with water. Half fill the spray tank with clean water and add the required amount of product. Agitate thoroughly before adding remainder of water. BOOM SPRAYING - Use 30-120L/ha of water. AERIAL SPRAYING - Use 10-90L/ha of water.

**EQUIPMENT MAINTENANCE AND USAGE**

Equipment that has been used for this chemical should not be used for the application of other materials to sensitive plants unless it has been well washed out with Tank & Equipment Cleaner followed by several clear water rinses.

**COMPATIBILITY/TANK MIXES**

This product can be mixed with Ally\*/Associate®, Kamba®, Bromicide®, Flowable Diuron, Flowable Igran\*, Glean\*/Lusta®, Glyphosate, Roundup PowerMAX\*, Grasp\*, Nugrass\*/Hoegrass\*, Nugran\*/Logran\*, Nugrex\*/Tigrex\*, Lontrel\*/Archer®, Propon, Puma\*. When tank mixing, add the required amount of compatible product when the spray tank is nearly full and while continuously agitating. If left overnight, agitate thoroughly before spraying. When mixing with Nugrass or Hoegrass, add an approved surfactant and DO NOT exceed 615mL of LVE Agritone per hectare. For consistent results, use a minimum of 80L water per ha by ground and 30L by air, and if wild oats are present, use 2L Nugrass per hectare.

**RESISTANT WEEDS WARNING**

GROUP	<b>I</b>	HERBICIDE
-------	----------	-----------

Nufarm LVE Agritone Selective Herbicide ("LVE Agritone") is a member of the Phenoxy group of herbicides. LVE Agritone has the Disruptors of plant cell growth mode of action. For weed resistance management LVE Agritone is a Group I herbicide. Some naturally-occurring weed biotypes resistant to LVE Agritone and other Group I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by LVE Agritone or other Group I herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Nufarm Australia Limited accepts no liability for any losses that may result from the failure of LVE Agritone to control resistant weeds.

**PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS**

DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. Avoid spray drift and vapour movement onto susceptible crops such as cotton, vines, tomatoes, vegetables, fruit trees, lucerne, legume crops and pastures, oilseed crops or other susceptible crops or trees (eg. kurrajongs, belahs, eucalypts).

**Legume Tolerance:** The tolerance of pasture legumes to this product can vary with species, variety, growing conditions, stage of growth and companion crop cover. While MCPA is preferable to 2,4-D in most situations, this product will cause greater damage than MCPA amine formulations. Lucerne and medics should not be sprayed under any circumstances with this product as severe damage will result.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT**

DO NOT contaminate streams, rivers or water ways with the chemical or used container

**STORAGE AND DISPOSAL (20L, 200L)**

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemical on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, triple rinse, break, crush or puncture and deliver empty packaging for appropriate disposal at an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Empty containers and product must not be burnt.

**(15L)**

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Triple or preferably pressure rinse inner bladder before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean packaging to recycler or designated collection point. If not recycling, deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

**Refillable Containers (500L & 1000L)**

Store in the closed, original container in a cool well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of sale for refill or storage.

**Envirodrum 110L Mini Bulk Returnable Container**

Store in original sealed Envirodrum in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the Envirodrum with water or any other foreign matter. After each use of the product, please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase. The Envirodrum remains the property of Nufarm Australia Limited.

**SAFETY DIRECTIONS**

Harmful if swallowed. Attacks eyes. Will irritate the skin. When preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms, and face thoroughly with soap and water. After each day's use wash gloves, goggles and contaminated clothing.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone: Australia 13 11 26).

**ADDITIONAL STATEMENTS (WHS REGULATIONS 2011)**

**Harmful in contact with skin. Harmful if inhaled. Causes serious eye irritation.** Avoid breathing spray. Wash hands and exposed skin thoroughly after handling. Do not eat, drink or smoke when using this product. Wear protective gloves, clothing, eye and face protection.

**SAFETY DATA SHEET**

For further information refer to the Safety Data Sheet (SDS), which can be obtained from your supplier or from the Nufarm website – [www.nufarm.com.au](http://www.nufarm.com.au)

**In case of emergency: Phone 1800 033 498** Ask for shift supervisor. Toll free 24 hours.

**CONDITIONS OF SALE**

"Any provisions or rights under the Competition and Consumer Act 2010 or relevant state legislation which cannot be excluded by those statutes or by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies, expressed or implied under common law, statute or otherwise, in relation to the sale, supply, use or application of this product, are excluded. Nufarm Australia Limited and/or its affiliates ("Nufarm") shall not accept any liability whatsoever (including consequential loss), or howsoever arising (including negligence) for any damage, injury or death connected with the sale, supply, use or application of this product except for liability which cannot be excluded by statute."

Nufarm Australia Limited  
ACN 004 377 780  
103-105 Pipe Road  
Laverton North Victoria 3026  
Tel: (03) 9282 1000  
Fax: (03) 9282 1001

© Archer, Associate, Bromicide, Kamba, Lusta, Nugrass and Nugrex are registered trade marks of, or used under licence by Nufarm Australia Limited.

\* Other registered trade marks

**WEED TABLE: Specific Weed Control Recommendations.**

Where weeds are to be sprayed in a crop or pasture, use only the rates given for the particular crop or situation indicated under the Directions of Use table. In most cases this will give control, however some hard to kill weeds or those in advanced stages of growth may only be suppressed. The rates listed in the weeds table below are for use where weeds only are present and crop or pasture is not involved. Use at these rates in a crop or pasture other than for spot spraying may cause damage.

WEEDS	RATE	CRITICAL COMMENTS
Bathurst burr ( <i>Xanthium spinosum</i> )	1.4L/ha	Moderately susceptible. Spray young seedlings.
Capeweed ( <i>Arctotheca calendula</i> )	1.49L/ha	Apply up to 2-4 leaf stage only when plants are no larger than 5cm in diameter.
Charlock ( <i>Sinapis arvensis</i> )	440mL-1.4L/ha	Young rosettes very susceptible.
Dandelion ( <i>Taraxacum officinale</i> )	1.4L/ha	Apply in early rosette stage.
Fat hen ( <i>Chenopodium album</i> )	880mL-1.4L/ha	Tolerance increases with age. Use lower rate only on seedlings.
Fumitory ( <i>Fumaria</i> )	965mL/ha	Moderately susceptible. Spray up to first true leaf stage. (Not Tas).
Hedge mustards ( <i>Sisymbrium</i> spp)	490-880mL/ha	Apply at rosette stage.
Lincoln weed ( <i>Diplotaxis tenuifolia</i> )	615mL/ha	Apply up to rosette stage only.
London rocket ( <i>Sisymbrium irio</i> )	965mL/ha	Spray at young stage.
Noogoora burr ( <i>Xanthium occidentale</i> )	1.4L/ha	Tolerance increases with age. Spray young seedlings only.
Paterson's curse ( <i>Echium plantagineum</i> )	1.49L/ha	Apply at 2 to 4 leaf stage only.
Saffron thistle ( <i>Carthamus lanatus</i> )	965mL-1.66L/ha	Spray young rosettes before leaves become spiny.
Scotch (cotton) thistle ( <i>Onopordum acanthium</i> )	615mL-1.49L/ha	Spray young rosettes.
Skeleton weed ( <i>Chondrilla juncea</i> )	965mL-1.4L/ha	Moderately susceptible. Spray rosettes before aerial growth commences.
Slender thistle ( <i>Carduus tenuiflorus</i> )	1.84L/ha	Moderately susceptible. Spray young rosettes.
Spear thistle ( <i>Cirsium vulgare</i> )	1.31-1.84L/ha	Moderately susceptible. Spray young rosettes.
Stinkwort ( <i>Dittrichia graveolens</i> )	1.4L/ha	Spray very young before plants 15cm high.
Turnip weed ( <i>Rapistrum rugosum</i> )	615-965mL/ha	Apply up to rosette stage. Apply lower rates to seedlings.
Volunteer canola ( <i>Brassica napus</i> )	1.31L/ha	Apply to seedlings only.
Variegated thistle ( <i>Silybum marianum</i> )	735-965mL/ha	Spray at pre-cabbage stage.
Volunteer safflower ( <i>Carthamus tinctorius</i> )	880mL/ha	Apply to seedlings only.
Volunteer sunflower ( <i>Helianthus annuus</i> )	1.31L/ha	Apply to seedlings only.
Wild radish ( <i>Raphanus raphanistrum</i> )	965mL-1.4L/ha	Apply up to 2 to 3 leaf stage only.
Wild sage ( <i>Salvia verbenaca</i> )	965mL/ha	Spray young seedlings.
Wild turnip ( <i>Brassica</i> spp)	440mL-1.31L/ha	Spray young rosettes. Lower rates for seedlings.