

Stewardship

24 Hour risk profile for Summer spraying

Always follow label instructions

Windspeeds must be above **4 km/h** and **less than 15-20 km/h** (refer to label) blowing away from sensitive areas

Medium spray quality: Delta T: 2-10

Coarse spray quality: Delta T: 2-12

To start spraying, the sun should be about 20 degrees above the horizon, and wind speed and direction consistent for 30-40 minutes

SUNRISE



Windspeed should be **above 4-5 km/h** after Sunrise to start spraying

20°



Spraying can only occur if the operator can be certain that a surface temperature inversion is not present.

The safest option is not to spray during this period.

Only use XC or UC spray quality, reduce spraying speed and boom height to minimise risk of droplets remaining airborne.

For more information, visit nufarm.com.au/stewardship

© 2018 - 2030 Nufarm Australia Ltd. All trade marks (®) are owned by Nufarm or used under license.

MIDDAY



Likely to be the best conditions for spraying



Be aware of higher evaporation risk and thermal activity after midday. Monitor plant stress.



Prepare to **STOP** all spraying if windspeeds start to drop



***EXTREME CAUTION REQUIRED**

High inversion risk
Dangerous air movement

Plan NOT TO SPRAY during this period



CAUTION REQUIRED

Surface Inversion onset likely.
Wind must be **above 11-12 km/h**



SUNSET



Often spraying into the early evening is possible in Summer when air movement has continued to mix the air and prevent a surface temperature inversion forming.



Pay very close attention to changes in wind speed and wind direction through out the evening.

Later in the evening air movement can become too unpredictable for safe spraying.

Often by 10-11 pm it has become unsafe.



MIDNIGHT

