



An suspension concentrate containing 100 g/L Cerevisane.  
A systemic resistance inducer against foliar fungal diseases for use in Oilseed Rape and Sugar Beet.

**IMPORTANT INFORMATION  
FOR USE ONLY AS A PROFESSIONAL FUNGICIDE**

Crop	Maximum Individual Dose (L product/ha)	Maximum Number of Applications	Latest Time of Application	Pre Harvest Interval
Sugar beet, spinach*, red beet*, spinach beet*, fodder beet*	2.0	5	BBCH 49 Root at harvestable size	1 day
Oilseed rape, flat-seed false flax*, gold-of-pleasure*, mustards*	2.0	1	BBCH 72 20% pods at final size	1 day

\* For these uses, growers are advised to test a small area first to establish crop safety.

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.**

To avoid risks to human health and the environment, comply with the instructions for use.

## SAFETY PRECAUTIONS

### OPERATOR PROTECTION

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

**WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND SUITABLE RESPIRATORY PROTECTIVE EQUIPMENT\*** when handling the concentrate or applying the product.

\* Disposable filtering facepiece respirator to at least EN149 FFP3 or equivalent.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

Micro-organisms may have the potential to provoke sensitising reactions.

### STORAGE AND DISPOSAL

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank and dispose of safely.

DO NOT RE-USE CONTAINER for any purpose.

### NUFARM UK LIMITED

Wyke Lane, Wyke, Bradford - West Yorkshire - BD12 9EJ - United Kingdom

Technical Helpline telephone number: +44 (0)1274 694714

24-hour emergency telephone number: +44 (0)1274 696603

PROTECT FROM FROST

SHAKE WELL BEFORE USE

FOR PROFESSIONAL USE ONLY

# 10 Litres e

## DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the product label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

### Restrictions

CONSULT PROCESSOR BEFORE USING ON SPINACH AND RED BEET GROWN FOR PROCESSING.

### Mode of action

Cerevisane is a host plant defence inducer (FRAC MOA code P06) so prepares plants for defense against fungal attacks. Apply preventatively, before the onset of disease.

### Resistance Management:

NUXINE contains Cerevisane A member of FRAC Code PO6, there are no reports of resistance to NUXINE, however alternating with fungicides of different modes of action is advisable for resistance management.

### Crop specific information

#### OILSEED RAPE, FLAT-SEED FALSE FLAX, GOLD-OF-PLEASURE, MUSTARDS

Pest	Root rot ( <i>Sclerotinia sclerotiorum</i> )
Application rate	2L product/ha
Maximum number of treatments	1
Notes and Timing:	Applications may be made between BBCH 65 (full flowering) and BBCH 72 (20% of pods at full size)
Level of control:	Reduction

#### SUGAR BEET, SPINACH, RED BEET, FODDER BEET, SPINACH BEET

Pest	Beet Leaf Spot ( <i>Cercospora beticola</i> )
Application rate	2L product/ha
Maximum number of treatments	5
Interval between treatments	14 days
Notes and Timing:	Applications may be made between BBCH 31 (Beginning of crop cover: leaves cover 10% of the ground) and BBCH 49 (Beet root has reached harvestable size)
Level of control:	Useful reduction. Levels of reduction can be variable and additional control measures may be required.

### Water volume

Apply in 200-300 L/ha water. Increase volume where crop is dense to ensure good coverage of foliage.

### Compatibility

NUXINE is compatible with a number of other products. Contact Nufarm UK Ltd for full details on compatibilities. Always read the label of the partner product carefully before use.

### Mixing

Half-fill the spray tank with clean water, add the required quantity of NUXINE and begin agitation. Maintain agitation until spraying is completed.

Rinse container thoroughly by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling.

### Spray tank clean-out

Clean spray tank hoses thoroughly before storing and using again.

## COMPANY ADVISORY INFORMATION

This section is not part of the Product Label under the Plant Protection Products Regulations 2011. It provides additional advice on use of the product at the discretion of the applicant.

## ACKNOWLEDGEMENTS

Nuxine is a trademark of Nufarm

### Conditions of supply

All goods supplied by us are of high grade and we believe them to be suitable but (as we cannot exercise any control over their mixing or use) all conditions and warranties, statutory or otherwise, as to the quality and fitness for any purpose of our goods are excluded, except in so far as such exclusion is prevented by law, and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use. This product is to be used only in accordance with the recommendations and instructions given on the labels provided with this pack. Use in any other circumstances is entirely at users risk.

To access the Safety Data Sheet for this product scan the QR code or refer to the Nufarm website at <https://www.nufarm.com/uk/>



An suspension concentrate containing 100 g/L Cerevisane.  
A systemic resistance inducer against foliar fungal diseases for use in Oilseed Rape and Sugar Beet.

**IMPORTANT INFORMATION  
FOR USE ONLY AS A PROFESSIONAL FUNGICIDE**

Crop	Maximum Individual Dose (L product/ha)	Maximum Number of Applications	Latest Time of Application	Pre Harvest Interval
Sugar beet, spinach*, red beet*, spinach beet*, fodder beet*	2.0	5	BBCH 49 Root at harvestable size	1 day
Oilseed rape, flat-seed false flax*, gold-of-pleasure*, mustards*	2.0	1	BBCH 72 20% pods at final size	1 day

\* For these uses, growers are advised to test a small area first to establish crop safety.

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.**

To avoid risks to human health and the environment, comply with the instructions for use.

**SAFETY PRECAUTIONS**

**OPERATOR PROTECTION**

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND SUITABLE RESPIRATORY PROTECTIVE EQUIPMENT\* when handling the concentrate or applying the product.

\* Disposable filtering facepiece respirator to at least EN149 FFP3 or equivalent.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

Micro-organisms may have the potential to provoke sensitising reactions.

**STORAGE AND DISPOSAL**

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank and dispose of safely.

DO NOT RE-USE CONTAINER for any purpose.

**NUFARM UK LIMITED**

Wyke Lane, Wyke, Bradford - West Yorkshire - BD12 9EJ - United Kingdom

Technical Helpline telephone number: +44 (0)1274 694714

24-hour emergency telephone number: +44 (0)1274 696603

PROTECT FROM FROST

SHAKE WELL BEFORE USE

FOR PROFESSIONAL USE ONLY

**10 Litres e**

510012771\_112024