



SELECTIVE HERBICIDE

TAKE DOWN TOUGH WEEDS WITH E-2.

Nothing knocks down unsightly weeds like E-2 herbicide. E-2 combines three powerful active ingredients to control more than 240 broadleaf weeds while providing safety for most desirable grasses and reducing the potential for resistant weed development.

COMPETITIVE ADVANTAGES

THE POWER OF THREE IN ONE FORMULATION

- Offers the power of three proven herbicides in one convenient pre-mixed formulation
- Excellent control of more than 240 broadleaf weeds including kochia, horsetail, field bindweed, thistles, plantain and other hard-to-control species
- Can be used for selective weed control in desirable grasses and as a tank-mix partner with products like ProClipse™ for cleaning up emerged weeds on bareground areas such as utility substations, industrial areas, highway medians and guardrails
- High-performing weed control in pasture and rangeland with control of tough species like hemp dogbane and dogfennel

AT-A-GLANCE INFO

EPA REG. NO. 228-442

ACTIVE INGR. 2,4-D (39.53%),
dicamba (4.10%),
fluroxypyr (5.90%)

FORMULATION liquid

CHEM. FAMILY phenoxyaliphatic acid,
pyridines, benzoic acids

GROUP NO. 4

MODE OF ACT. auxin growth regulator

SIGNAL WORD warning

RESTRICTED no

PACKAGE SIZE 1.5 gal container (5 per
case), 2.5 gal container
(2 per case), 250 gal tote

MORE PRODUCTS EXCLUSIVELY FROM NUFARM

Lock Down™ SC | **Scorch™ EXT** | **Razor™ Xtreme**

 **Nufarm**
Grow a better tomorrow



TAKE DOWN TOUGH WEEDS WITH E-2.

KEY USES

Rights-of-way, railroads and roadsides, industrial sites, fencerow and farmstead, non-irrigation ditch banks, recreational areas, pasture and rangeland, Conservation Reserve Programs (CRP) Land

KEY WEEDS CONTROLLED


(See product label for complete list)

Alder
Beggartick
Black-eyed Susan
Black medic
Buckhorn
Carolina geranium
Carpetweed
Chicory
Clover
Dock
False sunflower
Garlic mustard
Ground ivy
Hemp
Henbit
Honeysuckle
Jimsonweed
Kochia
Knotweed
Lambsquarters
Marshelder
Morningglory
Mouseear chickweed
Mustard
Pigweed
Purslane
Ragweed
Shepherd's-purse
Spatterdock
Speedwell
Spurge
Stinging nettle
Tansy ragwort
Thistle
Tumbleweed
Vetch
Virginia creeper
Wild four-o'clock
Willow
Yarrow
Yellow rocket

USE SUMMARY

WEED CONTROL PERCENTAGES

Field Bindweed Source: Dr. Phil Westra, Colorado State University, 2009
Buckhorn Plantain Source: P.L. Hipkins and H.L. Witt, Virginia Tech Noncrop and Aquatic Weed Research, 2009
Wild Carrot Source: P.L. Hipkins and H.L. Witt, Virginia Tech Noncrop and Aquatic Weed Research, 2009



NON-CROP

For control of broadleaf weeds, mix 2 to 5 pts/A in adequate water to thoroughly saturate all weeds with spray mixture. This may require a spray volume of 50 to 300 gals of water/A.

PASTURES, RANGELAND AND GRASS (HAY, SILAGE)

Some weed species will require tank mixes for adequate control. For these uses, mix 2 to 5 pts/A. Retreatment may be made as needed; however the maximum seasonal application rate is 2 lbs 2,4-D acid equivalent per acre per application. Minimum of 21 days between applications, limited to 2 applications per year. The pre-harvest interval (PHI) is 7 days (for cut forage for hay).

GRAZING AND FEEDING NON-LACTATING ANIMALS: There is no waiting period between treatment and grazing for non-lactating animals. Do not permit meat animals being finished for slaughter to graze treated fields within 30 days of slaughter.

GRAZING AND FEEDING LACTATING ANIMALS: Do not graze lactating dairy animals within 7 days of treatment.

DRY HAY AND SILAGE: Treated grasses may be harvested for dry hay or silage but do not harvest within 7 days of treatment.

CRP AND GENERAL FARMSTEAD

Some weed species require tank mixes for adequate control. For these uses, mix 2 to 5 pts/A. Retreatment may be made as needed; however the maximum seasonal application rate is 2 lbs 2,4-D acid equivalent per acre per application. For program lands, such as Conservation Reserve Program, consult program rules to determine whether grass or hay may be used. The more restrictive requirements of the program rules or this label must be followed.

See label for complete directions for use.



nufarm.com/usivm

Before applying any Nufarm product, confirm that it is registered for use in your state. For specific application rates, directions, mixing instructions and precautions, read the product label. Please visit www.nufarm.com/usivm to download a full label.

©2024 Nufarm. Important: Always read and follow label directions. Garlon® and Milestone® are trademarks of Corteva Agriscience.

15-VM-0008-D

