

Clash™

SELECTIVE HERBICIDE

STRIKE OUT TOUGH WEEDS.

Clash™ controls difficult ALS-, glyphosate- and triazine-resistant annual weeds in addition to a broad spectrum of perennial and biennial weeds. With its wide application window, Clash gives you the convenience and flexibility to deal with unpredictable spring weather making it well suited for early spring as well as fall applications when targeting difficult broadleaf weeds. Clash is labeled for a variety of sites from crop protection to vegetation management to provide use-flexibility no matter the market.

COMPETITIVE ADVANTAGES

A FLEXIBLE FORMULA FOR TOUGH WEED CONTROL

- Controls a broad spectrum of annual and perennial broadleaf weeds
- Ideal solution for ALS-, glyphosate- and triazine-resistant weeds
- Flexible application timing: preplant burndown, pre- and post-emergence in multiple crops
- Offers a wide application window when targeting broadleaf weeds in vegetation management sites

AT-A-GLANCE INFO

EPA REG. NO.	228-615
ACTIVE INGR.	DGA dicamba (56.8%) (4 lb ae/gal)
FORMULATION	liquid
CHEM. FAMILY	benzoic acid
GROUP NO.	4
MODE OF ACT.	auxin growth regulator
REI	24 hours
SIGNAL WORD	caution
RESTRICTED	no
PACKAGE SIZE	2.5 gal container (2 per case), 265 gal tote

MORE PRODUCTS EXCLUSIVELY FROM NUFARM



Piper™
HERBICIDE

Nufarm
Polaris AC Complete

 **Nufarm**
Grow a better tomorrow

Clash™

STRIKE OUT TOUGH WEEDS.

KEY USES

Selective broadleaf weed and brush control on non-crop lands: Rights-of-way (roadways, utility, railroad, highway, pipeline, and rights-of-way that run through pasture and rangeland), utility facilities (substations, pipelines, tankfarms, pumping stations, parking and storage areas, non-irrigated ditchbanks, and fencerows), fencerows, natural areas and forest site preparation

KEY WEEDS CONTROLLED

(See product label for complete list)

ANNUALS

Amaranth, Palmer
Broomweed, common
Carpetweed
Evening primrose, cutleaf
Kochia
Lambsquarters, common
Marestail (Horseweed)
Morningglory; ivyleaf, tall
Nightshade, black
Pennycress, field
Pigweed; prostrate, redroot
Ragweed; common, giant
Shepherd's-purse
Thistle, Russian
Waterhemp; common, tall
Waterprimrose, winged
Wormwood

BIENNIALS & PERENNIALS

Fern, bracken
Garlic, wild
Horsenettle, Carolina
Ivy, ground
Knapweed
Milkweed, common
Nettle, stinging
Nightshade, silverleaf
Plantain; bracted, broadleaf, buckhorn
Ragweed, western
Sericea lespedeza
Sowthistle
Spurge, leafy

WOODY

Kudzu
Russian olive
Spruce
Willow, witchhazel

USE SUMMARY

Clash™ can be applied to actively growing weeds as aerial, broadcast band, spray or cut surface treatments using water or sprayable fertilizer as a carrier. See label for complete instructions per application method.

ANNUALS

Small, actively growing: 8-16 fl oz/acre
Established weed growth: 16-24 fl oz/acre

BIENNIALS

Rosette (1-3" diameter): 8-16 fl oz/acre
Rosette (3" or larger diameter): 16-32 fl oz/acre
Bolting: 32 fl oz/acre

PERENNIALS

Top growth suppression: 8-16 fl oz/acre
Top growth control: 16-32 fl oz/acre

WOODY BRUSH AND VINES

Top growth suppression: 16-32 fl oz/acre
Top growth control: 32 fl oz/acre
Stems and stem suppression: 32 fl oz/acre

Approved surfactants, sprayable fertilizers or crop oil concentrate may be added to improve post-emergence weed control, particularly in dry growing conditions.

See product label for complete directions for use.

FOREST SITE PREPARATION

Clash may be used for control of undesirable woody species as well as many broadleaf weeds, vines, brambles, hardwood brush, and trees in forest site preparation. Clash may be applied as broadcast foliar sprays from ground or aerial equipment. Clash is absorbed through the leaf surfaces quickly after spraying and will also be absorbed from the soil by the roots. Translocation through the leaves, stems, and roots provides control of undesirable young conifer and broadleaf species. Woody plants, brush, and trees may not display the full extent of herbicide efficacy until several months following treatment. Clash provides application flexibility for extended windows of application and tank mix options (refer to Mixing and Application Procedures and Tank Mix Options).

APPLICATION INSTRUCTIONS

Ground operated spray equipment: Thoroughly mix and apply the specified amount of this product (2 pt per acre maximum) in a minimum of 15 gals of water per acre. Spray solution should uniformly cover undesirable foliage for best results.

See product label for complete directions for use.



nufarm.com/uscrop

For specific application rates, directions, mixing instructions and precautions, read the product label. Please visit www.nufarm.com/uscrop to download a full label.

©2020 Nufarm. Important: Always read and follow label instructions. Clash™ and Polaris® are trademarks of Nufarm. Piper® is a trademark of Valent U.S.A. LLC.



20-VM-0003