

Nufarm

Weedone® LV4 EC

SELECTIVE HERBICIDE

THE LOWEST VOLATILE 2,4-D ESTER ON THE MARKET.

Weedone® LV4 is the lowest volatile 2,4-D ester on the market. It is registered for small grains, corn, sorghum, pasture and rangeland, non-crop areas and preplant corn and soybeans. Weedone LV4 is more effective than amine formulations for controlling hard-to-kill weeds such as bindweed, curly dock, smartweeds, tansy ragwort, thistle and wild garlic.

COMPETITIVE ADVANTAGES

- Low volatile 2,4-D ester formulation
- Broad-spectrum control of broadleaf and woody weeds
- Extremely low freezing temperature (less than 0°F)
- Excellent compatibility with liquid fertilizers

AT-A-GLANCE INFO

EPA REG. NO.	228-139-71368
ACTIVE INGR.	2,4-D ester (67.2%)
FORMULATION	liquid
CHEM. FAMILY	phenoxy-carboxylic acid
GROUP NO.	4
MODE OF ACT.	synthetic auxin growth regulator
REI	12 hours
SIGNAL WORD	caution
RESTRICTED	no
PACKAGE SIZE	2.5 gal container (2 per case), 30 gal, 265 gal tote

MORE PRODUCTS EXCLUSIVELY FROM NUFARM

 BurnMaster

 Panther SC

 WeedMaster®

 Nufarm
Grow a better tomorrow

Nufarm
Weedone® LV4 EC

THE LOWEST VOLATILE 2,4-D ESTER ON THE MARKET.

KEY USES

Corn (field, sweet and popcorn)
Cereal grains
Grass seed crops
Non-crop areas (fencerows)
Pasture
Rangeland
Red Potatoes
Sorghum
Soybeans (preplant only)

KEY WEEDS CONTROLLED

(See product label for complete list)

Bindweed
Buckhorn
Canada thistle
Chicory
Cocklebur
Dock, curly
Dogfennel
Garlic, wild
Goldenrod
Ground ivy
Knotweed
Kochia
Lambsquarters
Locoweed
Marshelder
Morningglory
Mustard
Nettle
Nutgrass
Pigweed
Plantain
Poison ivy
Prickly lettuce
Primrose
Puncturevine
Ragweed
Russian thistle
Shepherd's-purse
Smartweeds
Tansy mustard
Thistle
Vervain

USE SUMMARY

For best results, apply this product as a water or oil spray during warm weather when young succulent weeds or brush are actively growing. Application under drought conditions often will give poor results.

The lower recommended rates will be satisfactory on susceptible, annual weeds. For perennial weeds and conditions such as the very dry areas of the Western States where control is difficult, the higher recommended rates should be used. Deep-rooted perennial weeds such as Canada thistle and field bindweed and many woody plants usually require repeated applications for maximum control.

Unless otherwise recommended, suggested application rates may be 1 to 10 gallons of total spray by air or 5 to 25 gallons by ground application equipment. If band treatment is used, base the dosage rate on the actual area to be sprayed. Although water quantities may vary due to different types of application equipment, sufficient water must be used to provide for complete and uniform coverage. Higher water gallonage may be used if desired to improve spray coverage. In all cases, use the same recommended amount of 2,4-D per acre.

When product is used for weed control in crops, the growth stage of the crop must be considered. For crop uses, do not mix with oil or other adjuvants unless specifically recommended on label. To do so may reduce herbicide's selectivity and could result in crop damage.

See product label for complete directions for use.



nufarm.com/uscrop

For specific application rates, directions, mixing instructions and precautions, read the product label. Please visit www.nufarm.com/uscrop to download a full label.

©2021 Nufarm. Important: Always read and follow label instructions. BurnMaster™, Panther®, Weedone® and WeedMaster® are trademarks of Nufarm.

