

Candor™

SELECTIVE HERBICIDE

CONTROL TREES AND BRUSH WITH A LOW VOLATILE PREMIX.

Candor™ herbicide can be used to control weeds in pastures and rangeland, Conservation Reserve Program (CRP) acres, non-irrigation ditchbanks, fence rows, roadsides and other non-crop areas and industrial sites. The post-emergence activity of Candor targets woody plants and brush – including blackberries and poison ivy – as well as annual and perennial broadleaves, while leaving desirable grasses unharmed.

COMPETITIVE ADVANTAGES

TOUGH WEED CONTROL

- Controls more than 100 tough annual and perennial broadleaf weed and woody plants without harming desirable grasses
- Proven active ingredients in a pre-mix formulation

FLEXIBLE APPLICATION

- May be combined with liquid nitrogen fertilizer
- Can be applied by ground equipment, helicopter or by spot treatment
- No grazing restrictions for most livestock, including cattle, horses, gestating mares, sheep and goats
- Rainfast in 2 hours
- Non-restricted use pesticide

AT-A-GLANCE INFO

EPA REG. NO.	228-565
ACTIVE INGR.	triclopyr (16.5%), 2,4-D (34.4%)
FORMULATION	liquid
CHEM. FAMILY	pyridinecarboxylic acid
HRAC NO.	4
MODE OF ACT.	synthetic auxin
REI	24 hours
SIGNAL WORD	caution
RESTRICTED	no
PACKAGE SIZE	2.5 gal container (2 per case)

MORE PRODUCTS EXCLUSIVELY FROM NUFARM

 BurnMaster

 Cheetah

 Panther Pro


Nufarm
Grow a better tomorrow

CONTROL TREES AND BRUSH WITH A LOW VOLATILE PREMIX.

KEY USES

Conservation Reserve Program (CRP) land, fallow systems, general farmstead, pastures, rangeland, fencerows, non-irrigation ditchbanks, roadsides, other non-crop areas and industrial sites

KEY WEEDS CONTROLLED

(See product label for complete list)

Amaranth, spiny
Blackberry
Burdock
Buttercup
Chickweed, mouseear
Clover, white sweet
Clover, bur
Cocklebur
Croton, woolly
Dandelion
Dock, curly
Dogbane, hemp
Galinsoga, hairy
Goatsbeard
Henbit
Horseweed
Ironweed, tall, western
Ivy, ground
Kochia
Kudzu
Lambsquarters
Lespedeza
Lettuce, wild
Mustard
Oxalis
Pennycress, field
Pepperweed, field
Pigweed, redroot
Plantain
Poison ivy
Purslane, annual
Ragweed
Radish, wild
Ragwort, tansy
Shepherd's-purse
Sneezeweed, bitter
Sowthistle, annual
Spurge
Sunflower
Thistle
Vetch
Violet, wild

USE SUMMARY

Candor™ can be applied in-season to foliage or during dormancy. For best results from foliar applications, apply when foliage is actively growing.

Rates range from 1 quart to 2 gallons per acre and vary depending on target weeds and timing of applications.

See product label for complete directions for use.

GRAZING AND HAYING RESTRICTIONS

GRAZING AND HARVESTING GREEN FORAGE

- No grazing restrictions for livestock including cattle, horses (including gestating mares), sheep, goats, etc.
- Lactating dairy animals — Do not graze or harvest green forage from treated area for 14 days after treatment.

HAYING (HARVESTING OF DRIED FORAGE)

- Livestock: Do not harvest hay for 7 days after treatment.
- Lactating dairy animals: Do not harvest hay until the next growing season.

See product label for complete grazing and haying restrictions.

SLAUGHTER RESTRICTIONS

Withdraw livestock from grazing treated grass or consumption of treated hay at least 3 days before slaughter. This restriction applies to grazing during the season of treatment or hay harvested during the season of treatment.

See product label for slaughter restrictions.

