

This seed has been treated with SATIVA IMF SEMBOLITE MAX. When treated at the recommended use rate of 3.4 – 5.0 oz/100lbs of seed, this seed has been treated with 0.026 – 0.039 lbs imidacloprid, 0.0013 – 0.002 lbs Metalaxyl, 0.001 – 0.0016 lbs Tebuconazole, and 0.0008 – 0.0012 lbs fludioxonil per 100 lbs of seed.

CAUTION **TREATED SEED**

DO NOT USE FOR FEED, FOOD, OR OIL PURPOSE. STORE AWAY FROM FEEDS AND FOODSTUFFS. DO NOT ALLOW CHILDREN, PETS OR LIVESTOCK TO HAVE ACCESS TO TREATED SEED.

PRECAUTIONARY INFORMATION FOR TREATED SEED

Seed Treated with SATIVA IMF MAX is harmful if swallowed.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Wear long-sleeved shirt, long pants, and chemical resistant gloves when handling treated seed.

Wash hands before eating, drinking, and chewing gum, using tobacco or using the toilet. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Pollinator Precautions: Imidacloprid is highly toxic to bees. Ensure that planting equipment is functioning properly in accordance with manufacturer specifications to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.

Groundwater Advisory Statement: Metalaxyl is known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

Exposed treated seed may be hazardous to birds. Cover or incorporate spilled treated seeds. Excess or leftover seed may be double-sown around the headland or buried away from bodies of water in accordance with local requirements.

Treated seed must be planted into the soil at a depth greater than 1 inch.

USE RESTRICTIONS:

The maximum application rate (including seed treatment, foliar application, and soil application) per acre per year for imidacloprid is 0.5 lbs.

DO NOT graze or feed livestock on treated areas for 45 days after planting.

REPLANTING / CROP ROTATION RESTRICTIONS:

In the event of a crop failure or harvest of a crop grown from SATIVA IMF SEMBOLITE MAX treated seed, the field may be replanted immediately to artichoke, barley, borage, Brassica (cole) leafy vegetables, bulb vegetables, canola, cilantro, corn (field or sweet), cotton, cranberry, crambe, cucurbits, eggplant, flax, groundcherry, leafy petiole vegetables, leafy vegetables, legume vegetables (succulent or dried including soybean), millet, mustard seed, oats, okra, pepinos, pepper, popcorn, potato, rapeseed, rye, safflower, sorghum, soybean, strawberry, sugarbeets, sunflower, tomatillo, tomato, triticale, root and tuber vegetables, watercress, and wheat. For cereals including buckwheat, and rice; the minimum plant-back interval is 30 days from the date SATIVA IMF SEMBOLITE MAX treated seed was planted. Cover crops for soil building or erosion control may be planted at any time; but do not graze or harvest for food or feed. For all other crops not listed on an imidacloprid label, or for crops for which no imidacloprid, tolerance for the active ingredient has been established, a 12-month plant-back interval must be observed.

DISPOSAL:

Dispose of seed packaging in accordance with local requirements.

DO NOT contaminate water bodies when disposing of planting equipment wash water.

Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticides remain in the ethanol by-products that are used in agronomic practice.

**For Chemical Spill, Leak, Fire, or Exposure,
Call CHEMTREC (800) 424-9300**

**For Medical Emergencies Only,
Call (877) 325-1840**

RV 052516
13845000

Sativa is a Registered Trademark of Nufarm Americas Inc.

Nufarm Americas Inc.
AGT Division
11901 S. Austin Ave.
Alsip, IL 60803

