

This seed has been treated with fludioxonil and metalaxyl fungicides at the rates specified by the manufacturer and imidacloprid insecticide at a rate of 0.0625 lbs imidacloprid/ 100 lbs of seed.

CAUTION

TREATED SEED

DO NOT USE TREATED SEED FOR FEED, FOOD, OR OIL PURPOSES. STORE AWAY FROM FEEDS AND FOODSTUFFS. DO NOT ALLOW CHILDREN, PETS OR LIVESTOCK TO HAVE ACCESS TO TREATED SEEDS. DO NOT USE THE SAME AUGER FOR TREATED SEED THAT WILL HANDLE COMMERCIAL OR FEED GRAINS.

PRECAUTIONARY INFORMATION FOR TREATED SEED

For treated seed: May be harmful if swallowed. Eye contact may cause moderate irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Wear long pants, long-sleeved shirt, shoes plus socks and chemical resistant gloves made of any waterproof material when opening this bag or handling (e.g., loading, pouring) treated seed.

After treated seed has been planted, **DO NOT** enter or allow worker entry into areas planted with treated seeds during the restricted-entry interval (REI) of 24 Hours. **EXCEPTION:** Workers may enter these areas without restriction if there will be no worker contact with the soil/media subsurface.

RESTRICTIONS:

Treated seed must be planted into the soil at a depth greater than 1 inch.

DO NOT harvest mustard greens. Rape greens grown and harvested from metalaxyl fungicide treated seed must not be used for human and feed consumption. Rapeseed grown and harvested from metalaxyl fungicide-treated seed is only for industrial uses and cannot be used for edible oil or any other human/feed consumption.

DO NOT use metalaxyl or mefenoxam for control of Pythium in water-seeded rice.

The maximum application rate (including seed treatment, foliar application, and soil application) per acre per year for imidacloprid is 0.5 lbs. Forage may not be grazed until 30 days after planting.

REPLANTING / CROP ROTATION RESTRICTIONS:

In the event of a crop failure or harvest of a crop grown from SENATOR 600 FS (imidacloprid) treated seed, the field may be replanted immediately to artichoke, barley, borage, Brassica (cole) leafy vegetables, bulb vegetables, canola, cilantro, corn (field or sweet), cotton, cranberry, crambe, cucurbits, eggplant, flax, groundcherry, leafy petiole vegetables, leafy vegetables, legume vegetables (succulent or dried including soybean), millet, mustard

seed, oats, okra, pepinos, pepper, popcorn, potato, rapeseed, rye, safflower, sorghum, soybean, strawberry, sugarbeets, sunflower, tomatillo, tomato, triticale, root and tuber vegetables, watercress, and wheat. For cereals including buckwheat, and rice; the minimum plant-back interval is 30 days from the date SENATOR 600 FS (imidacloprid) treated seed was planted. Cover crops for soil building or erosion control may be planted at any time; but do not graze or harvest for food or feed. For all other crops not listed on an imidacloprid label, or for crops for which no imidacloprid, tolerance for the active ingredient has been established, a 12-month plant-back interval must be observed.

ENVIRONMENTAL HAZARDS

Exposed treated seed on the soil surface may be hazardous to birds and other wildlife. Cover or collect treated seed spilled during loading and planting. Excess or leftover seed may be double-sown around the headland or buried away from bodies of water in accordance with local requirements.

DO NOT contaminate bodies of water when disposing of planting equipment washwater. If excess treated seed remains after planting, dispose of it by burying away from bodies of water.

Pollinator Precautions: Imidacloprid is highly toxic to bees. Ensure that planting equipment is functioning properly in accordance with manufacturer specifications to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.

SEED STORAGE:

Store treated seed in cool, dry conditions. **DO NOT** store leftover seed where there is a chance of it becoming mixed with untreated seed.

SEED DISPOSAL:

Dispose of seed packaging in accordance with local requirements. **DO NOT** contaminate water bodies when disposing of planting equipment wash water.

Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticides remain in the ethanol by-products that are used in agronomic practice.

**For Chemical Spill, Leak, Fire, or Exposure,
Call CHEMTREC (800) 424-9300**

**For Medical Emergencies Only,
Call (877) 325-1840**

RV 081015[1]
13713000

Nufarm Americas Inc.
AGT Division
11901 S. Austin Ave.
Alsip, IL 60803

