

This seed has been treated with fludioxonil and metalaxyl fungicides at the rates specified by the manufacturer.

CAUTION

TREATED SEED

DO NOT USE FOR FEED, FOOD, OR OIL PURPOSES. STORE TREATED SEED AWAY FROM FEEDS AND FOODSTUFFS. DO NOT ALLOW CHILDREN, PETS OR LIVESTOCK TO HAVE ACCESS TO TREATED SEEDS. DO NOT USE THE SAME AUGER FOR TREATING SEED THAT WILL HANDLE COMMERCIAL OR FEED GRAINS.

PRECAUTIONARY INFORMATION FOR TREATED SEED

For treated seed: May be harmful if swallowed. Eye contact may cause moderate irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Wear long pants, long-sleeved shirt, shoes plus socks and chemical resistant gloves made of any waterproof material when opening this bag or handling (e.g., loading, pouring) treated seed.

After treated seed has been planted, **DO NOT** enter or allow worker entry into areas planted with treated seeds during the restricted-entry interval (REI) of 24 Hours. **EXCEPTION:** Workers may enter these areas without restriction if there will be no worker contact with the soil/media subsurface.

RESTRICTIONS:

Forage may not be grazed until 30 days after planting.

DO NOT plant any crop not listed on this label within 30 days of planting metalaxyl fungicide-treated seed.

DO NOT harvest mustard greens. Rape greens grown and harvested from metalaxyl fungicide treated seed must not be used for human and feed consumption. Rapeseed grown and harvested from metalaxyl fungicide-treated seed is only for industrial uses

and cannot be used for edible oil or any other human/feed consumption.

DO NOT use metalaxyl or mefenoxam for control of *Pythium* in water-seeded rice.

ENVIRONMENTAL HAZARDS

Exposed treated seed on the soil surface may be hazardous to birds and other wildlife. Cover or collect treated seed spilled during loading and planting. Excess or leftover seed may be double-sown around the headland or buried away from bodies of water in accordance with local requirements.

DO NOT contaminate bodies of water when disposing of planting equipment washwater. If excess treated seed remains after planting, dispose of it by burying away from bodies of water.

SEED STORAGE:

Store treated seed in cool, dry conditions. **DO NOT** store leftover seed where there is a chance of it becoming mixed with untreated seed.

SEED DISPOSAL:

Dispose of seed packaging in accordance with local requirements.

DO NOT contaminate water bodies when disposing of planting equipment wash water.

Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticides remain in the ethanol by-products that are used in agronomic practice.

**For Chemical Spill, Leak, Fire, or Exposure,
Call CHEMTREC (800) 424-9300**

**For Medical Emergencies Only,
Call (877) 325-1840**

RV 110812

13711000

Nufarm Americas
Inc. AGT Division
11901 S. Austin
Ave. Alsip, IL 60803

