

This seed has been treated with Tebuconazole (SATIVA 309 FS) and Metalaxyl (SEBRING 480 FS) fungicide at the rates specified by the manufacturer.

# CAUTION

## TREATED SEED

**DO NOT USE FOR FEED, FOOD, OR OIL PURPOSES. KEEP OUT OF REACH OF CHILDREN. STORE TREATED SEED AWAY FROM FEEDS AND FOODSTUFFS. DO NOT ALLOW CHILDREN, PETS OR LIVESTOCK TO HAVE ACCESS TO TREATED SEEDS. DO NOT USE THE SAME AUGER FOR TREATING SEED THAT WILL HANDLE COMMERCIAL OR FEED GRAINS.**

### PRECAUTIONARY INFORMATION FOR TREATED SEED

The treatment applied to this seed may be harmful if swallowed, absorbed through the skin, or inhaled. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

### PERSONAL PROTECTIVE EQUIPMENT (PPE):

Wear long pants, long-sleeved shirt, shoes plus socks, and chemical-resistant gloves made of any waterproof material when opening this bag or handling (e.g., loading, pouring) treated seed.

Wash hands before eating, drinking, and chewing gum, using tobacco or using the toilet. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

After treated seed has been planted, **DO NOT** enter or allow worker entry into areas planted with treated seeds during the restricted-entry interval (REI) of 24 hours. **EXCEPTION:** Workers may enter these areas without restriction if there will be no worker contact with the soil/media subsurface.

### ENVIRONMENTAL HAZARDS

Treated seed exposed on soil surfaces may be hazardous to wildlife. Cover or collect treated seed spilled during loading and planting.

## **USE RESTRICTIONS:**

**IMPORTANT:** Barley, wheat, and oat green forage may be grazed or harvested 31 days after seeding. All other forage may not be grazed until 30 days after planting. Do not plant any crop not listed on the Sebring 480 FS label within 30 days of planting Sebring 480 FS treated seed. Do not harvest mustard greens. Rape greens grown and harvested from Sebring 480 FS Fungicide-treated seed must not be used for human and feed consumption. Rapeseed grown and harvested from Sebring 480 FS Fungicide-treated seed is only for industrial uses and cannot be used for edible oil or any other human/feed consumption.

## **DISPOSAL:**

**DO NOT** contaminate bodies of water when disposing of planting equipment washwater. If excess treated seed remains after planting, dispose of it by burying away from bodies of water.

**DO NOT** store leftover treated seed where there is a chance of it becoming mixed with untreated seed.

Store treated seed in cool, dry conditions.

Dispose of treated seed packaging in accordance with local requirements.

Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticides remain in the ethanol by-products that are used in agronomic practice.

**For Chemical Spill, Leak, Fire, or Exposure,  
Call CHEMTREC (800) 424-9300**

**For Medical Emergencies Only,  
Call (877) 325-1840**

RV042016

**13776000**

Sativa and Sebring are Registered Trademarks of Nufarm Americas Inc.

Nufarm Americas Inc.  
11901 S. Austin Ave.  
Alsip, IL 60803

