

This seed has been treated with Difenoconazole (SALIENT 372 FS) seed treatment fungicide.

CAUTION

TREATED SEED

DO NOT USE FOR FEED, FOOD, OR OIL PROCESSING. KEEP OUT OF REACH OF CHILDREN. STORE AWAY FROM FEEDS AND OTHER FOODSTUFFS. DO NOT ALLOW CHILDREN, PETS OR LIVESTOCK TO HAVE ACCESS TO TREATED SEEDS.

PRECAUTIONARY INFORMATION FOR TREATED SEED

The treatment applied to this seed may be harmful if swallowed, absorbed through the skin, or inhaled. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Wear long pants, long-sleeved shirt, shoes, socks, and chemical-resistant gloves when opening this bag or loading/pouring the treated seed.

After treated seed has been planted, do not enter or allow worker entry into areas planted with treated seeds during the restricted-entry interval (REI) of 12 hours. Exception: If the treated seed is soil-injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: coveralls, chemical-resistant gloves made of any waterproof material, and chemical-resistant footwear.

ENVIRONMENTAL HAZARDS

Cover, incorporate, or collect treated seed spilled during loading and planting.

USE RESTRICTIONS:

Green wheat, oats, rye or triticale forage may not be grazed until 55 days after planting. Green barley forage may not be grazed until 30 days after planting. Do not plant any crop other than those on this label within 30 days to fields in which treated seeds were planted.

DISPOSAL:

DO NOT contaminate bodies of water when disposing of planting equipment washwater. If excess treated seed remains after planting, dispose of it by burying away from bodies of water.

DO NOT store leftover treated seed where there is a chance of it becoming mixed with untreated seed.

Store treated seed in cool, dry conditions.

Dispose of treated seed packaging in accordance with local requirements.

Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticides remain in the ethanol by-products that are used in agronomic practice.

**For Chemical Spill, Leak, Fire, or Exposure,
Call CHEMTREC (800) 424-9300
For Medical Emergencies Only,
Call (877) 325-1840**

(RV102716)

13867000

Salient is a Registered Trademark of Nufarm Americas Inc.

Nufarm Americas Inc.
AGT Division
11901 S. Austin Ave.
Alsip, IL 60803

