

SPIRATO IMT^m 348 FS

This seed has been treated with (imidacloprid, metalaxyl, thiophanate-methyl, and fludioxonil).

CAUTION TREATED SEED

This seed has been treated with

0.0625	lbs	_____	mg	Imidacloprid
0.0156	lbs	_____	mg	Metalaxyl
0.00118	lbs	_____	mg	Thiophanate-methyl
0.0025	lbs	_____	mg	Fludioxonil
X	Per 100	LBS OF SEED		
	Per Seed			

DO NOT USE FOR FEED, FOOD, OR OIL PURPOSES. STORE AWAY FROM FEEDS AND FOODSTUFFS. DO NOT ALLOW CHILDREN, PETS OR LIVESTOCK TO HAVE ACCESS TO TREATED SEED.

PRECAUTIONARY INFORMATION FOR TREATED SEED

Seed Treated with SPIRATO IMT^m 348 FS is harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Store treated seeds away from feeds and foodstuffs.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Note: Persons handling product, treating seed, or involved in bagging treated seed, sewing or moving bags of treated seed, or cleaning up bagging areas or seed treatment equipment must wear the following PPE: long-sleeved shirt, long pants and chemical-resistant gloves when handling treated seed.

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Groundwater Advisory Statement: Metalaxyl is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Pollinator Precautions: Imidacloprid is highly toxic to bees. Ensure that planting equipment is functioning properly in accordance with manufacturer specifications to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.

Treated seed must be planted into the soil at a depth greater than 1 inch.

Exposed treated seed may be hazardous to birds and wildlife. Dispose of all excess treated seed and seed packaging or containers by burial away from bodies of water in accordance with any local requirements. Cover or incorporate spilled treated seeds. **DO NOT** contaminate bodies of water when disposing of planting equipment wash water.

RESTRICTIONS:

The maximum application rate (including seed treatment, foliar application, and soil application) per acre per calendar year for imidacloprid is 0.5 lbs.

DO NOT graze or feed livestock on treated areas for 45 days after planting.

DO NOT apply a neonicotinoid insecticide within 45 days of planting seed treated with SPIRATO IMTm 348 FS.

In the event of a crop failure or harvest of a crop grown from SPIRATO IMTm 348 FS treated seed, the field may be replanted immediately to artichoke, barley, borage, Brassica (cole) leafy vegetables, bulb vegetables, canola, cilantro, corn (field or sweet), cotton, cranberry, crambe, cucurbits, eggplant, flax, groundcherry, leafy petiole vegetables, leafy vegetables, legume vegetables (succulent or dried including soybean), millet, mustard seed, oats, okra, pepinos, pepper, popcorn, potato, rapeseed, rye, safflower, sorghum, soybean, strawberry, sugarbeets, sunflower, tomatillo, tomato, triticale, root and tuber vegetables, watercress, and wheat. For cereals including buckwheat, and rice; the minimum plant-back interval is 30 days from the date SPIRATO IMTm 348 FS treated seed was planted. Cover crops for soil building or erosion control may be planted at any time; but do not graze or harvest for food or feed. For all other crops not listed on an imidacloprid label, or for crops for which no imidacloprid tolerance for the active ingredient has been established, a 12-month plant-back interval must be observed.

DISPOSAL:

Dispose of seed packaging in accordance with local requirements. Excess or leftover seed may be double-sown around the headland or buried away from bodies of water in accordance with local requirements. **DO NOT** contaminate water bodies when disposing of planting equipment wash water.

Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticides remain in the ethanol by-products that are used in agronomic practice.

**For Chemical Spill, Leak, Fire, or Exposure,
Call CHEMTREC (800) 424-9300**

**For Medical Emergencies Only,
Call (877) 325-1840**

RV 102116

13714000

Nufarm Americas Inc.
AGT Division
11901 S. Austin Ave.
Alsip, IL 60803

