



SUPPLEMENTAL LABEL

Nufarm T-Methyl 4.5 F Fungicide

EPA Reg. No. 228-652
01-05-2012

LABEL REVISIONS

Use Directions for Pre-Harvest Use to Control Post-Harvest Disease on Apples
Use Directions for Beans, Dry and Succulent, & PHI for Lima Beans
Use Directions for Suppression of Crown Rot on Strawberries

READ AND FOLLOW THE ENTIRE LABEL BOOKLET FOR THIS PRODUCT BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING. "Label" as used in this supplemental labeling refers to the label booklet for this product and this supplemental.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN ANY MANNER INCONSISTENT WITH ITS LABELING.
This label must be in the possession of the user at the time of application.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS IN THE PRODUCT LABEL BOOKLET MUST BE FOLLOWED, INCLUDING STATEMENTS PERTAINING TO THE WORKER PROTECTION STANDARDS, ON THE EPA REGISTERED LABEL.

APPLES

Pre-Harvest Use to Control Post-Harvest Diseases

DISEASES	SCIENTIFIC NAME	FLUID OZ/ACRE	FLUID OZ/100 GAL	USE INFORMATION
Storage rot blue mold Gray mold Bull's eye rot	<i>Penicillium expansum</i> <i>Botrytis cinerea</i> <i>Neofabraea</i> spp.	1 (0.035 lb ai/A)	3.75 - 5	Apply as a pre-harvest spray within 2 weeks to 3 days of harvest. Thorough coverage of the fruit is required. Application closer to harvest may provide better efficacy. For resistance management, do not use a benzimidazole fungicide (i.e. Mertect®) post-harvest following a pre-harvest application of a product containing thiophanate-methyl. Application of a non- benzimidazole post-harvest fungicide will provide additional protection from post-harvest diseases.

NOTES:

- ♦ Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 2 days.
- ♦ Do not apply more than 80 fl. oz. of product (2.8 lb ai) per acre per season.
- ♦ Pre-harvest Interval (PHI): 24 hours

BEANS, DRY AND SUCCULENT

Including: Lima bean, Snap bean, Kidney bean, Mung bean, Navy bean, Pinto bean, Wax bean, Broad bean, Fava bean, Asparagus bean, Blackeyed pea, Cowpea, Sweet lupine, White lupine, White Sweet lupine, Grain lupine, Chick pea, Garbanzo bean

DISEASES	SCIENTIFIC NAME	FLUID OZ/ACRE	USE INFORMATION
Gray mold White mold Anthracnose	<i>Botrytis</i> sp. <i>Sclerotinia</i> sp. <i>Colletotrichum</i>	30 - 40 (1.0 - 1.4 lb ai/A)	For one application: Apply when 100% of plants have at least one open bloom or when conditions are favorable for disease development.
		20 - 30 (0.7 - 1.0 lb ai/A)	For multiple applications: Make first application when 10%-30% of plants have at least one open bloom. Follow with additional applications at 4- to 7-day intervals. Apply prior to the development of disease for best results.
NOTES:			
<ul style="list-style-type: none"> ♦ For succulent beans, do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. ♦ For dry beans, do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 3 days. ♦ Do not apply more than 80 fl. oz. of product (2.8 lb ai) per acre per crop cycle. ♦ Pre-harvest Interval (PHI) – <ul style="list-style-type: none"> <u>All States Except California:</u> 14 days for succulent beans and lima beans 28 days for dry beans <u>California only:</u> 14 days for succulent beans 28 days for lima beans and dry beans 			

STRAWBERRIES

DISEASES	SCIENTIFIC NAME	FLUID OZ/ACRE	USE INFORMATION
Crown rot * (Suppression only)	<i>Colletotrichum</i> spp.	15 - 20 (0.53 - 0.7 lb ai/A)	Begin applications after establishment of the transplants and continue through bloom at 10 to 14 day intervals. Use the high rate if the fields have a history of <i>Colletotrichum</i> crown rot and/or conditions are favorable for disease development. Will not control <i>Phytophthora</i> species.
NOTES:			
<ul style="list-style-type: none"> ♦ Do not apply more than 80 fl. oz. product (2.8 lb ai) per acre per year. ♦ Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. ♦ Pre-harvest Interval (PHI): 1 day <p>* Not registered for use in California</p>			

NOTICE

Read the "WARRANTY DISCLAIMER" and "LIMITATION OF LIABILITY" in the label booklet for this product before using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

**Manufactured For NUFARM AMERICAS INC.
150 Harvester Drive, Burr Ridge, IL 60527**