

AS-50®

Agricultural Streptomycin

This seed has been treated with AS-50® (Streptomycin) to control halo blight.

DO NOT use for food, feed, or oil purposes. Store away from feeds and other foodstuffs.

CAUTION

TREATED SEED

**DO NOT USE TREATED SEED FOR
FEED, FOOD, OR OIL PURPOSES.**

KEEP OUT OF REACH OF CHILDREN

PRECAUTIONARY INFORMATION FOR TREATED SEED

DO NOT allow children, pets or livestock to have access to treated seeds.

Eye exposure to the seed treatment may cause moderate irritation. Harmful if swallowed or inhaled. Harmful if absorbed through skin. Avoid breathing dust from treated seed. Avoid contact with eyes, skin, or clothing.

Wear long pants, long-sleeved shirt, shoes, socks, and chemical-resistant gloves when opening this bag or loading/pouring the treated seed.

After the seeds have been planted, do not enter or allow worker entry into treated areas during the **restricted-entry interval (REI)** of **12 hours**. Exception: If the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: coveralls, chemical-resistant gloves made of any waterproof material, and chemical-resistant footwear.

ENVIRONMENTAL HAZARDS

Exposed treated seed on the soil may be hazardous to wildlife. Dispose of all excess treated seed and seed packaging or containers by burial away from bodies of water in accordance with any local requirements. Cover, incorporate, or collect treated seeds spilled during loading and planting. **DO NOT** contaminate bodies of water when disposing of planting equipment wash water.

DISPOSAL:

Dispose of seed packaging in accordance with local requirements.

DO NOT contaminate water bodies when disposing of planting equipment wash water.

Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticides remain in the ethanol by-products that are used in agronomic practice.

**For Chemical Spill, Leak, Fire, or Exposure,
Call CHEMTREC (800) 424-9300**

**For Medical Emergencies Only,
Call (877) 325-1840**

RV 081110

13743000

Nufarm Americas Inc.
AGT Division
150 Harvest Drive
Burr Ridge, IL 60527

