

This seed has been treated with Fludioxonil (SPIRATO 480 FS), Metalaxyl (SEBRING 480 FS), and Thiophanate-methyl (ST-METHYL 540 FS) fungicides at the rates specified by the manufacturer.

CAUTION

TREATED SEED

DO NOT USE TREATED SEED FOR FEED, FOOD, OR OIL PURPOSES. STORE AWAY FROM FEEDS AND FOODSTUFFS. DO NOT ALLOW CHILDREN, PETS OR LIVESTOCK TO HAVE ACCESS TO TREATED SEEDS.

PRECAUTIONARY INFORMATION FOR TREATED SEED

Harmful if swallowed or inhaled. Causes moderate eye irritation. Harmful if absorbed through the skin. Avoid breathing dust from treated seed. Avoid contact with eyes, skin, or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Wear long-sleeved shirt, long pants, shoes, socks, and chemical resistant gloves when handling treated seed.

Wash hands before eating, drinking, and chewing gum, using tobacco or using the toilet. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Exposed treated seed may be hazardous to birds and wildlife. Dispose of all excess treated seed and seed packaging or containers by burial away from bodies of water in accordance with any local requirements. Cover, incorporate, or collect treated seeds spilled during loading and planting. Do not contaminate bodies of water when disposing of planting equipment wash water. Excess or leftover seed may be double-sown around the headland or buried away from bodies of water in accordance with local requirements.

Groundwater Advisory Statement: Metalaxyl is known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

USE RESTRICTIONS:

After the seeds have been planted, do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. Exception: If the treated seed is soil-injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: coveralls, chemical-resistant gloves made of any waterproof material, and chemical-resistant footwear.

DO NOT graze or feed livestock on treated areas for 45 days after planting.

DISPOSAL:

Dispose of seed packaging in accordance with local requirements.

DO NOT contaminate water bodies when disposing of planting equipment wash water.

Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticides remain in the ethanol by-products that are used in agronomic practice.

**For Chemical Spill, Leak, Fire, or Exposure,
Call CHEMTREC (800) 424-9300**

**For Medical Emergencies Only,
Call (877) 325-1840**

RV011513

13712000

Spirato and Sebring are Registered Trademarks of Nufarm Americas Inc.

Nufarm Americas Inc.
AGT Division
150 Harvest Drive
Burr Ridge, IL 60527

