

Comet™

ECONOMICAL POST-EMERGENT CONTROL OF BROADLEAF WEEDS.

Comet™ is a selective post-emergence herbicide that controls annual and perennial broadleaf weeds and woodybrush in rangeland and permanent pastures. Comet may also be used for selective post-emergence control of annual and perennial broadleaf weeds and volunteer potatoes in small grains, field corn, sweet corn, grain sorghum, fallow cropland, on-farm non-cropland and pome fruit.

Comet may be applied as a single broadcast treatment or spot treatment to control susceptible broadleaf weeds in on-farm non-cropland areas such as fence rows, building perimeters, around irrigation equipment and on-farm private roadways.

COMPETITIVE ADVANTAGES

- Excellent control of broadleaf weeds
- Can be tank-mixed with other broadleaf and grass herbicides as well as most fungicides, insecticides and foliar fertilizers
- Offers rotational flexibility to next year's crop

AT-A-GLANCE INFO

EPA REG. NO.	71368-87
ACTIVE INGR.	fluroxypyr (26.2%)
FORMULATION	liquid
CHEM. FAMILY	carboxylic acid
HRAC NO.	4
MODE OF ACT.	synthetic auxin growth regulator
REI	12 hours
SIGNAL WORD	warning
RESTRICTED	no
PACKAGE SIZE	2.5 gal container (2 per case), 265 gal tote

MORE PRODUCTS EXCLUSIVELY FROM NUFARM





ECONOMICAL POST-EMERGENT CONTROL OF BROADLEAF WEEDS.

KEY USES

Rangeland
Permanent pastures
Small grains, field corn, sweet corn,
grain sorghum, fallow cropland
and on-farm non-cropland
Pome fruit

KEY WEEDS CONTROLLED

(See product label for complete list)

Bedstraw (cleavers)
Bindweed, field*
Bindweed, hedge
Buckwheat, wild*
Canola, volunteer*
Chickweed
Clover, white
Cocklebur
Coffeeweed
Devilsclaw*
Flax, volunteer
Grape spp.
Hemp dogbane
Horsetail, field*
Horseweed (marestail)*
Jimsonweed
Knotweed*
Kochia*
Mallow, common*
Mallow, Venice
Marshelder*
Morningglory
Mustard spp.*
Nightshade spp.*
Pennycress, field*
Potato, volunteer*
Prickly lettuce
Puncturevine
Purslane, common
Ragweed, common
Sunflower
Thistle, Russian*
Velvetleaf

* Suppression

USE SUMMARY

Avoid unintentional spray drift to adjacent susceptible broadleaf plants or broadleaf crops.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Use application rates of 2/3 to 1 1/3 pints per acre depending on crop (2/3 to 2 2/3 for pome fruit). One application is allowed per season.

See label for complete directions for use.



nufarm.com/uscrop

For specific application rates, directions, mixing instructions and precautions, read the product label. Please visit www.nufarm.com/uscrop to download a full label.

©2020 Nufarm. Important: Always read and follow label instructions. BurnMaster™, Comet™ and Patriot® are trademarks of Nufarm.



15-AG-0019-E